


**EFFECT OF DIFFERENT RATES OF NITROGEN AND
PHOSPHOROUS APPLICATION ON THE GROWTH AND YIELD
OF BARI TOMATO-2 (RATAN) (*Lycopersicon esculentum*. Mill)**

BY

MD. MANJURUL ALAM

Registration No: 08-03200



A Thesis

submitted to the Department of Soil Science,
Sher-e-Bangla Agricultural University, Dhaka
in partial fulfillment of the requirement for the degree of

**MASTER OF SCIENCE (MS)
IN
SOIL SCIENCE**

Semester: July-December, 2009

Approved By:

Mohammad Mosharraf Hossain
Supervisor
Assistant Professor
Department of Soil Science
Sher-e-Bangla Agricultural University
Dhaka- 1207

Dr. Gopi Nath Chandra Sutradhar
Co-supervisor
Professor
Department of Soil Science
Sher-e-Bangla Agricultural University
Dhaka- 1207

Dr. Md. Asaduzzaman Khan
Associate Professor
Chairman &
Examination Committee
Department of Soil Science
Sher-e-Bangla Agricultural University
Dhaka- 1207





DEPARTMENT OF SOIL SCIENCE
Sher-e-Bangla Agricultural University
Sher-e-Bangla Nagar, Dhaka-1207

CERTIFICATE

This is to certify that the thesis entitled, **“EFFECT OF DIFFERENT RATES OF NITROGEN AND PHOSPHOROUS APPLICATION ON THE GROWTH AND YIELD OF BARI TOMATO-2 (RATAN) (*Lycopersicon esculentum*. Mill)”** submitted to the Faculty of Agriculture, Sher-e-Bangla Agricultural University, Dhaka, in partial fulfillment of the requirement for the degree of **MASTER OF SCIENCE IN SOIL SCIENCE**, embodies the result of a piece of *bona fide* research work carried out by **MD. MANJURUL ALAM**, Registration No. **08-03200** under my supervision and my guidance. No part of the thesis has been submitted for any other degree or diploma.

I further certify that any help or source of information, received during the course of this investigation has been duly acknowledged by him.

Date: 07-6-2011
Dhaka, Bangladesh

Mohammad Mosharraf Hossain
Supervisor
Assistant Professor
Department of Soil Science
Sher-e-Bangla Agricultural University
Dhaka-1207



Dedication

I dedicate my thesis to my parents whose efforts and ever willing support have made this dream come true

ACKNOWLEDGEMENT

The author expresses the endless gratitude to the supreme ruler Almighty Allah Who enabled him to complete the thesis successfully.

The author expresses his deepest sense of gratitude, immense indebtedness and profound appreciation to his reverend research supervisor **Mohammad Mosharraf Hossain**, Assistant Professor, Department of Soil Science, Sher-e-Bangla Agricultural University, Dhaka-1207 for planning the research work, vigilant supervision, constructive suggestions, sympathetic encouragement to conduct the research work as well as preparation and for going through the manuscript of the thesis.

He also expresses his grateful appreciation and deep sense of respect to his research co-supervisor **Dr. Gopi Nath Chandra Sutradhar**, Professor, Department of Soil Science, Sher-e-Bangla Agricultural University, Dhaka-1207 for proper guidance, continuous advice, constructive criticism, painstaking suggestions, kind help and encouragement during the course of research work and preparation of this manuscript.

He is immensely indebted to Associate Professor Dr. Md. Asaduzzaman Khan, Chairman of the Department of Soil Science, Sher-e-Bangla Agricultural University, Dhaka-1207 for providing him the facilities to conduct the experiment.

He expresses his gratitude and profound respect to Professor Dr. Md. Nurul Islam, Professor Dr. Alok Kumar Paul, Associate Professor Mr. A.T.M. Shamsuddoha, Associate Professor Dr. Md. Asaduzzaman Khan, Associate Professor Mst. Afrose Jahan, and all of his respective teachers of the Department of Soil Science, Sher-e-Bangla Agricultural University, Dhaka-1207 for their valuable teaching and co-operations in completing the research study.



He also expresses his sincere gratitude and thanks to all other staff of the Department of Soil Science, Sher-e-Bangla Agricultural University, Dhaka-1207 for their kind and untiring help and sincere cooperation during the research period.

He also expresses his thanks to Dr. Md. Manjurul Alam, Senior Scientific Officer, Bangladesh Institute of Nuclear Agriculture (BINA), Mymensingh for sympathetic help and co-operation for data analysis and preparation of this manuscript.

It is beyond words to express his appreciation, gratitude and esteemed respect for his parents for their unconditional love, ever willing help, patience, constant encouragement and their faith in him which always kept him focused on his objectives and helped to achieve his goals.

At last but not the least, the author wishes to express his profound gratitude and deepest appreciation to his all other relatives for their ever ending prayer, encouragement, sacrifice and dedicated efforts to educate him to this level.

Dhaka, Bangladesh
December 2009

The author

ABSTRACT

The experiment was conducted at the Research Farm of Sher-e-Bangla Agricultural University, Dhaka-1207, during November 2008 to March 2009 to evaluate the effect of nitrogen (N) and phosphorous (P) on the growth and yield component of BARI tomato-2 (Ratan), is one of the high yielding varieties of indeterminate type. The experiment was laid out in a Randomized Complete Block Design (RCBD) with three replications of each fertilizers treatment combination. The treatments used were: four levels of nitrogen as N_0 : 0 kg N/ha, N_{100} : 100 kg N/ha, N_{150} : 150 kg N/ha and N_{200} : 200 kg N/ha and four levels of phosphorous as P_0 : 0 kg P_2O_5 /ha, P_{50} : 50 kg P_2O_5 /ha, P_{100} : 100 kg P_2O_5 /ha and P_{150} : 150 kg P_2O_5 /ha with 16 treatment combinations. Yield and yield contributing characters, N and P concentration in plant and post harvest soil were recorded. Plant height at 60 days after transplanting, number of branches per plant, number of flowers per plant, number of fruits of plant and fruit weight per plant increased significantly with increasing N level up to 150 kg N/ha, whereas fruit yield increased significantly up to 200 kg N/ha. All these parameters also increased significantly with increasing the level of P up to 150 kg P/ha. However, the treatment combination $N_{200}P_{150}$ showed the maximum plant height at harvest (93.62 cm). On the other hand, the treatment combination $N_{200}P_{100}$ produced the maximum number of flowers per plant (49.86), number of fruits per cluster (3.99), number of fruits per plant (38.30) fruit weight (2.5 kg/plant) and fruit yield (89.93 t/ha). N and P concentrations in shoot and fruit were the highest in $N_{200}P_{150}$ treatment. The treatment had no significant effect on pH of post harvest soil. However increasing the application of nitrogen increased soil total N and P content up to N_{150} treatment. Application of P had not significant effect on total nitrogen but P_{150} produced the highest P content in soil. $N_{200}P_{150}$ combination had the highest N (0.119%) and P (44.77 ppm) concentration in plant.

LIST OF CONTENTS

CHAPTER	TITLE	PAGE NO.
	ACKNOWLEDGEMENT	i-ii
	ABSTRACT	iii
	LIST OF CONTENTS	iv-v
	LIST OF TABLES	vi-vii
	LIST OF FIGURES	viii
	LIST OF APPENDICES	ix
	LIST OF ACRONYMS	x-xii
1	INTRODUCTION	1-3
2	REVIEW OF LITERATURE	4-18
	2.1 Effect of nitrogen on the growth and yield of tomato	4
	2.2 Effect of Phosphorous on the growth and yield of tomato	12
	2.3 Integrated effect of nitrogen and phosphorous on the growth and yield of tomato	14
3	MATERIALS AND METHODS	19-32
	3.1 Experimental site	19
	3.2 Characteristics of soil	19
	3.3 Climate	21
	3.4 Seeds and variety	21
	3.5 Raising of seedlings	21
	3.6 Treatments	22
	3.7 Design and layout of experiment	23
	3.8 Collection and preparation of initial soil sample	23
	3.9 Land preparation	23
	3.10 Application of manure and fertilizers	25
	3.11 Transplanting of seedlings	25
	3.12 Irrigation and drainage	25
	3.13 Gap filling and weeding	25
	3.14 Pest management	26
	3.15 Harvesting	26
	3.16 Collection of experimental data	26
	3.17 Analytical Methods of Soil and Plant Sample Analysis	29
	3.18 Chemical analysis of plant samples	29
	3.19 Soil Sample Analysis	30
	3.20 Statistical Analysis	32

LIST OF CONTENTS (CONT'D)

CHAPTER	TITLE	PAGE NO.
4	RESULTS AND DISCUSSION	33-64
4.1	Yield and yield contributing character of tomato	33
4.1.1	Plant height	33
4.1.2	No. of leaves per plant at 60 DAT	38
4.1.3	No. of branches per plant at final harvest	39
4.1.4	No. of flower clusters per plant	42
4.1.5	No of flowers per cluster	43
4.1.6	No of flowers per plant	43
4.1.7	Number of fruits per cluster	44
4.1.8	Number of fruits per plant	45
4.1.9	Weight of individual fruit	46
4.1.10	Fruit size (Fruit length)	47
4.1.11	Fruit size (Fruit breadth)	48
4.1.12	Fruit weight per plant (kg)	49
4.1.13	Fruit yield (ton/ha)	52
4.2	Nutrient content in shoot	54
4.2.1	Nitrogen content (%) in shoot	54
4.2.2	Phosphorous content (ppm) in shoot	55
4.3	Nutrient content (%) in fruit	58
4.3.1	Nitrogen content (%) in fruit	58
4.3.2	Phosphorous content (%) in fruit	59
4.4	Status of post harvest soil	60
4.4.1	P ^H at post harvest soil	60
4.4.2	Nitrogen (%) at post harvest soil	60
4.4.3	Phosphorous content (ppm) at post harvest soil	64
5	SUMMERY AND CONCLUSION	65-66
	REFERENCES	67-75
	APPENDICES	76-79



LIST OF TABLES

SL. NO.	TITLE	PAGE NO.
1	Morphological characteristics of experimental field	20
2	Physical and chemical properties of the experimental initial soil	20
3	Different methods for determination of physico-chemical properties of soil Physical properties of soil.	31
4	Effect of nitrogen and phosphorous on plant height, number of leaves per plant and branches per plant of tomato	36
5	Interaction effect of nitrogen and phosphorous on plant height, number of leaves per plant and branches per plant of tomato	37
6	Effect of nitrogen and phosphorous on number of cluster / plant, number of flowers/ cluster, number of flowers/ plant, number of fruits/ cluster and number of fruits/ plant of tomato	40
7	Interaction effect of nitrogen and phosphorous on number of flower cluster/ plant, number of flowers/ cluster, number of flowers/ plant, number of fruits/ cluster and number of fruits/ plant of tomato	41
8	Effect of nitrogen and phosphorous on fruit size, weight of individual fruit, fruit weight/ plant and fruit yield of tomato	50
9	Interaction effect of nitrogen and phosphorous fruit size, weight of individual fruit (gm), fruit weight /plant (kg), fruit yield (ton/ha) of tomato.	51

LIST OF TABLES (CONT'D)

SL. NO.	TITLE	PAGE NO.
10	Effect of nitrogen and phosphorous on Nitrogen (%) and phosphorous (ppm) in shoot & N (%) and P (%) in fruit of tomato	56
11	Interaction effect of nitrogen and phosphorous on nitrogen (%) in shoot, phosphorous (ppm) in shoot , N % in fruit, P % in fruit of tomato	57
12	Effect of nitrogen and phosphorous on P ^H of post harvest soil, nitrogen (%), phosphorous (ppm) soil of post harvest soil	62
13	Interaction effect of nitrogen and phosphorous on pH of post harvest soil, nitrogen (%), phosphorous (ppm) soil of post harvest soil	63

LIST OF FIGURES

SI. NO.	TITLE	PAGE NO.
1	Field layout of the experimental plot	24
2	Effect of nitrogen on the yield (ton/ha) of tomato	53
3	Effect of phosphorous on the yield (ton/ha) of tomato	53
4	Interaction effect of nitrogen and phosphorous on the yield (ton/ha) of tomato	53



LIST OF APPENDICES

APPENDICES	TITLE	PAGE NO.
I	Map showing the experimental site under the study	76
II	Monthly record of air temperature, relative humidity and sunshine of the experimental site during the period from October 2008 to February 2009	77
III	Analysis of variance of the data on plant height an 30,60 DAT, at harvest, number of leaves per plant and 60DAT and number of branches per plant at harvesting period as influenced by nitrogen and phosphorous	77
IV	Analysis of variance of the data on number of cluster / plant, number of flowers/ cluster, number of flowers/ plant, number of fruits/ cluster, number of fruits/ plant, influenced by nitrogen and phosphorous	78
V	Analysis of variance of the data on fruit size, weight of individual fruit (gm), fruit weight /plant (kg) and fruit yield (ton/ha) as influenced by nitrogen and phosphorous	78
VI	Analysis of variance of the data on pH of post harvest soil, nitrogen (%), phosphorous (ppm) of post harvest soil as influenced by nitrogen and phosphorous	79
VII	Analysis of variance of the data on Nitrogen (%) in straw, Phosphorous (P)in ppm in straw , Potassium (K) in (%)straw, N% in fruit, P% in fruit, K% in fruit as influenced by nitrogen and phosphorous	79

LIST OF ACRONYMS

ABBREVIATIONS	ELABORATIONS
%	Percent
⁰ C	Degree Centigrade
AEZ	Agro-Ecological Zone
Anon.	Anonymous
ANOVA	Analysis of Variance
BARC	Bangladesh Agriculture Research Council
BARI	Bangladesh Agricultural Research Institute
BAU	Bangladesh Agricultural University
BBS	Bangladesh Bureau of Statistics
C	Carbon
CAN	Calcium Ammonium Nitrate
cc	Cubic Centimeter
cm	Centimeter
Conc.	Concentration
cv	Cultivar(s)
CV	Coefficient of Variance
DAP	Di Ammonium Phosphate
DAS	Days After Sowing
DAT	Days After Transplanting
DMRT	Duncan's Multiple Range Test
e.g.	Example
<i>et al.</i>	And other
etc	etcetera
FAO	Food and Agriculture Organization
Fed	feddon
gm	Gram

LIST OF ACRONYMS (CONT'D)


ABBREVIATIONS	ELABORATIONS
ha	Hectare
HCl	Hydrochloric acid
hr	Hour
HRC	Horticulture Research Centre
HYVs	High Yielding Varieties
Inst.	Institute
IRRI	International Rice Research Institute
<i>J.</i>	Journal
K	Potassium
kg	Kilogram
Kg/ha	Kilogram per hectare
KSI	Potassium Supplementation Index
LSD	Least Significant Difference
m	Meter
m ²	Square meter
ml	Milliliter
mm	Millimeter
Mn	Manganese
MOP	Muriate of Potash
N	Nitrogen
NaCl	Sodium Chloride
NAR	Net Assimilation Rate
NFT	Nutrient Film Technique
NGOs	Non Government Organizations
No.	Number
NS	Non Significant

LIST OF ACRONYMS (CONT'D)

ABBREVIATIONS

ELABORATIONS

NSI	Nitrogen Supplementation Index
P	Phosphorous
pH	Hydrogen ion concentration
ppm	parts per million
PSI	Phosphorous Supplementation Index
RCBD	Randomized Complete Block Design
Res.	Research
RH	Relative Humidity
RUE	Radiation Use Efficiency
SAU	Sher-E-Bangla Agricultural University
SRDI	Soil Research Development Institute
t/ha	Ton per hectare
TSP	Triple Superphosphate
TSS	Total Soluble Solid
UNDP	United Nations Development Program
USDA	United States Department of Agriculture
VAM	Vesicular Arbuscular Mycorrhiza
<i>viz.</i>	<i>videlicet</i>
Wt.	Weight



Chapter I

Introduction

CHAPTER I

INTRODUCTION

Tomato (*Lycopersicon esculentum*. Mill) belongs to the family solanaceae, is one of the most important, popular and nutritious vegetable in Bangladesh. It is originated in tropical America (Salunkhe *et al.* 1987) and ranks third in terms of world, vegetable production (FAO, 1997). It is the top listed canned vegetables (Choudhury, 1997). It is adapted to a wide range of climate ranging from tropics to within a few degrees of the Arctic Circle. However, in spite of its broad adaptation, it is concentrated in a few area and rather dry area (Cuortero and Fernandez, 1999).

The popularity of tomato and different products of tomato processing is increasing day by day. It is a nutritious and delicious vegetable used in salads, soups and processed into stable products like ketchup, sauce and juice etc. They are extensively used in the Canning food nutritive value of the fruit is an important aspect of quality in tomato and public demand. Food value of tomato is very rich because of high contents of vitamins A, B and C including calcium and carotene. Tomato adds flavor to the foods and it is also rich in medicinal value. In other aspect, Bangladesh can earn huge foreign currencies to export raw tomatoes and processed tomatoes by producing more quality tomatoes using modern technologies and high yielding varieties. For this purpose we should take initiatives to make our farmers more knowledgeable and expert to grow more food from our limited lands. In addition, we should encourage the farmers and other NGOs to grow more and more vegetable like tomato. The government should take the steps to establish tomato processing plant because of our huge production in all the year.

The edaphic and climatic conditions of winter seasons of Bangladesh are congenial for tomato cultivation. Very recently some variety was developed for year round cultivation of tomato. Among the winter vegetable crops in

Bangladesh, tomato ranks second in respect of production and third in respect of area (BBS, 2006). During the year 2005-2006, 1,31,000 metric tons of tomato were produced from 13,845 hectares of land with an average yield of 9.46 t/ha (BBS, 2006). Except this yield is quite low as compared to other leading tomato producing countries of the world such as China, Egypt, USA, Turkey, where per hectare yield was reported as 30.39, 34.00, and 41.77 t/ha, respectively.

The lower yield of tomato in Bangladesh, however, is not an incidence of the low yield potentiality of this crop, but, the fact that the lower yield may be attributed to a number of reasons viz. unavailability of quality seeds of improved varieties, fertilizers management, disease infestation and improper moisture management. Among them fertilizer management is a vital factor that influence the growth and yield of tomato. Balance fertilizations in crops will act as an insurance against possible nutrient deficiencies that may be created by the respected use of a single nutrient (Manage *et al.*, 1982). Among different nutrients that were required for tomato cultivation nitrogen and phosphorous are the most important nutrients. On the other hand, soils of Bangladesh have been deficient in nitrogen and phosphorous fertilizer. So, it is necessary to apply these nutrient elements for satisfactory growth, development and also yield of tomato (BARC, 2005). Tomato required large quantity of readily available fertilizer nutrients (Gupta and Shukla, 1977). In indeterminate type of tomato, vegetative and reproduction stages overlap and the plants need nutrients up to fruit ripening to get one ton fresh fruit, plants need to absorb, on average 2.5-3.0 Kg N, 0.2-0.3 Kg K (Hedge, 1997).

In Bangladesh, fertilizer especially nitrogen is the most critical input for increasing production and had appropriately been recognized as the central element for agricultural development (Mukhopadhyay *et al.*, 1986). More than any other nutrient, nitrogen influences vegetative growth and yield of tomato plant. Nitrogen is essential for building up of protoplasm and protein which induced cell division and initial meristematic activity when applies in optimum quantity (Sing and Kuer, 1969). Nitrogen had the largest effect on yield and quality of tomato (Xin *et al.*, 1997). It also promotes vegetative growth, flower and fruit setting of tomato

(Lose and Som, 1990). Optimum nitrogen increases fruit quality, fruit size, color, taste and acidity (Sharma and Mann, 1971). It significantly increases the growth and yield of tomato (Banerjee *et al.*, 1997).

Next to nitrogen fertilizer, phosphatic fertilizers dwell is the second most important input for increasing crop production. High level of phosphorous throughout root zone is essential for rapid root development and for good utilization of water and other nutrients by the plant. Phosphorous has profound effect on number of flowers that progressively increases the yield (Al-Afifi *et al.*, 1993, Razia and Islam, 1980) and marketable yield of tomato (Candilo, 1993).


The relationship between nitrogen and phosphorous to the tomato has received considerable attention and appears to have had a profound influence on horticultural practices. Combined effects of nitrogen and phosphorous have also been proved quite effective in increasing growth and yield of tomato (Kuksal *et al.*, 1977). Optimum rates of nitrogen and phosphorous not only increase the yield but also increase the quality of tomato (Sharma and Mann, 1971).

In Bangladesh, BARI (Bangladesh Agricultural Research Institute) has released at least ten HYVs (High yielding varieties) of tomato. The variety BARI Tomato-2 (Raton) is one of the high yielding varieties of indeterminate type. It may be cultivated throughout the year all over the country. Research in response of tomato to nitrogen, phosphorous, potassium and sulphur fertilization is very limited for tomato cultivation.

Keeping all the above facts in view, the present experiment has been undertaken with following objectives:

1. To find out the optimum doses of nitrogen and phosphorous for growth and yield of tomato.
2. To find out the combined effect of nitrogen and phosphorous for attaining desirable tomato yield.





Chapter II

Review of Literature

CHAPTER II

REVIEW OF LITERATURE

Tomato is one of the most important vegetable crops in Bangladesh and as well as many countries of the world. It is a high yielding year round crop and fertilizer responsive crop. Among different fertilizers, nitrogen and phosphorous are two of the essential plant nutrient elements and each plays a significant role in tomato production. But a very few studies in relation to nitrogen and phosphorous on growth, yield and development of tomato. The research work so far done in Bangladesh and is not adequate, specific and conclusive. Nevertheless, some of the important and informative works and research findings related to nitrogen and phosphorous fertilizers so far been done at home and abroad on this crop and have been reviewed in this chapter under the following parameters.

2.1 Effect of nitrogen on the growth and yield of tomato

Sainju *et al.* (2001) conducted an experiment at Agricultural Research Station, Fort Valley State University, Fort Valley to evaluated hairy vetch residue as nitrogen fertilizer for tomato in soilless medium. The ability of hairy vetch (*Vicia villosa* Roth) residue (100gm/plant) to supply N to increase yields of tomato (*Lycopersicon esculentum*) was compared with that of N fertilization (0, 4.1, and 8.2g N/plant) in a medium containing a mixture of 3 perlite : 1 vermiculite in a greenhouse. Leaf dry weight, leaf and stem N uptake of tomato, and NH_4^+ and inorganic N concentrations in the medium at transplanting were significantly greater than without residue. Nitrogen fertilization increased fruit number, fresh and dry yields and N uptake, stem, leaf, and root dry weights and N uptake, root length, and total dry weight and N uptake. The residue was as effective in increasing fresh fruit yield, total dry weight, and N uptake was 4.4 to 7.9 g/plant of N fertilizer. Tomato yield and N uptake per unit amount of N supplied was greater for the residue than N fertilization.

Research was conducted by Dufault *et al.* (2000) in Charleston, South Carolina, to determine (1) The effect of supplemental nitrogen (N) at 60 or 120 kg ha⁻¹ following winter cover crop of wheat, Tomato (*Lycopersicon esculentum mill.*) and snap bean (*Phaseolus vulgaris L*) grown in rotation; and (2) The distribution and retention of soil nitrates in the soil profile as affected by N fertilization and cover cropping. Total marketable tomato yield increased as fertilizer N increased to 60 kg ha⁻¹ in two out of four years and with 120 kg ha⁻¹ in one out of four years. In all cover crop or fallow plots, as fertilizer N application levels increased, the soil nitrates also increased.

Rhoads *et al.* (1999) carried out an experiment to evaluate the influence of N rates and ground cover following tomato on soil nitrate-N movement was monitored in spring and autumn crops grown at the Florida A&M University, Florida, USA. Nitrogen rates varied from 0 to 360 lb/acre in the spring crop and from 0 to 600 lb/acre in autumn tomato. Yield ranged from 1900 to 2600 boxes/acre in spring tomato, and from 1300 to 2700 boxes/acre in autumn tomato. Fertilizer N rates above 180 lb/acre were excessive, as shown by yield and residual soil nitrate N levels. Residual soil nitrate N was proportional to N application rate. Soil nitrate N concentration following harvest was highest in the 1 to 3 ft depth range for spring tomato and the 2 to 4 ft depth range for autumn tomato.

A study was conducted by Ceylan *et al.* (2001) at Odemis, Izmir, Turkey to assess the effect of ammonium nitrate and urea fertilizers at 0, 12, 24, 36 kg N /ha on nitrogen uptake and accumulation in tomato plants. The total nitrogen, NO₂-N and NO₃-N contents of leaches and fruits were determined. On the first and second harvest dates, the highest NO₃⁻ N and NO₂-N amounts in tomato leaches and fruits were obtained upon treatment with 36 kg N/ha. Ammonium nitrate application increased nitrate and nitrite accumulation compared to urea application. The highest was recorded upon treatment with 24 kg N.

An experiment was conducted in Uttar Pradesh, India, by Singh *et al.* (2000) to determine the suitable rate and application of N fertilizers for obtaining optimum growth and yield of tomato cv. Pusa Hybrid -2. N was applied at 40 kg/ha basal, 40kg/ha top dressing, 80 kg/ha in 2 splits (40 kg/ha basal+ 40 kg/ha top dressing), 50 kg/ha in 2 splits (40 kg/ha basal +10 kg/ha foliar), 60 kg/ha (40 kg/ha basal+ 20 kg/ha foliar), 70 kg/ha (40 kg/ha basal +30 kg/ha foliar) and 80 kg/ha (40 kg/ha basal +20 kg/ha top dressing +20 kg/ha foliar). N @ 80 kg/ha applied in 3 splits produced the highest yield and biomass. Increasing N rates resulted in increasing biomass and yield.

Two field experiments were conducted in Egypt by Awad *et al.* (2001) to study the effect of intercropping parsley and demsisa with tomato under 4 rates of N fertilizer (100, 120, 140, and 160 kg N /fed). The results showed that increasing N fertilizer rate enhanced total yield and net assimilation rate (NAR) of both mono and mixed crops, earliness index of tomato and NPK uptake of tomato in NAR, total yield, earliness index and N uptake. The best values were obtained by pure stand planting at the highest N rate (160 kg N/fed), whereas the best P and K uptake were attained at 140 and 120 kg N/fed, respectively. The highest value of N supplementation index (NSI) for tomato was obtained at 100 kg N /fed, whereas the highest values of phosphorous supplementation index (PSI) and potassium supplementation index (KSI) were recorded by plants which received 160 kg N/fed.

A field experiment was conducted by Manoj and Raghav (2001) to evaluating two F₁ hybrids of tomato, three plant spacing (75cm ×50cm, 75cm ×75cm, and 75cm ×100cm) and five levels of nitrogen (0,75, 150, 225, and 300 kg/ha) was conducted during 1995-96 and 1996-97 at the Research Station, Nagina of G.B. Pant University of Agriculture and Technology, Pantnagar (Uttar Pradesh, India) on sandy loam soil. Among the various levels of nitrogen, 300 kg/ha was found to be best in improving the growth and yield.

Sandoval *et al.* (2001) conducted an experiment to determine the tomato yield response to 0:100, 10:90, 20:80, 30:70, 40:60 $\text{NH}_4^+ \text{N} : \text{NO}_3^- \text{N}$ ratios supplied at the vegetative, vegetative plus flowering, flowering plus fruiting, and fruiting stages, and over the entire plant life cycle. Neither the length of $\text{NH}_4^+ \text{N}$ supply nor the $\text{NH}_4^+ \text{N}$ concentration in solution affected tomato yield. Plant height was not affected by $\text{NH}_4^+ \text{N}$ concentration in either the winter or spring experiments, and neither was fruit firmness measured for fruit at the mature green stage. Fresh and dry weights were unaffected by $\text{NH}_4^+ \text{N}$ concentration.

Bot *et al.* (2001) carried out an experiment to evaluate the response of adult tomato plants growing in rock wool in a greenhouse to N withdrawal from the nutrient solution was studied over a 6 week period during fruit production. The major effect of N withdrawal included the impairment of growth of fast growing organs. Fruit growth was impaired, leading to a reduction in yield. The growth of young leaves was also inhibited. The stores of nitrate N were depleted after removal of N in the solution, but it took 45 days for the plants to metabolize completely their nitrate reserves.

Tomato cv. Pusa Gaurav was treated with N @ 0, 40, 80 and 120 kg/ha K @ 0, 30 and 60 kg/ha Gupta and Sengar (2001) conducted an experiment in Madhya Pradesh, India during rabi 1992-93 and 1993-94. N application resulted in increases in plant height, number of fruits per plant, fruit weight and fresh yield. Increasing N rate produced a corresponding increase in yield components, except total soluble solids (TSS) content. K increased vegetative growth, yield and TSS content.

Sainju *et al.* (2000) conducted an experiment on cover crops to examine the influence of legume and N fertilizer application (0, 90, and 180 kg N/ha) on short and long-term effects on soil C and N ratio and tomato yield and N uptake. N uptake similar to that produced by 90 and 180 kg N/ha. Nitrogen fertilizer application increased PNM and inorganic N after split application, tomato yield and N uptake but decreased organic C and N and PCM.

Hoffland *et al.* (2000) conducted an experiment to study how nitrogen availability affects within plant allocation to growth and secondary metabolites. Tomato plants were grown at six levels of nitrogen availability. When nitrogen availability increased, plant relative growth rate increased, but tissue carbon/nitrogen ratio in the second oldest true leaf and allocation to large glandular trichomes as well as to the defense compounds rutin, chlorogenic acid decreased but leaf protein concentration increased.

Chang *et al.* (2001) conducted an experiment to investigate the effect of nitrogen supply by NH_4^+ deposit fertilizer on plant growth and nitrogen uptake of tomatoes. NH_4^+ deposit fertilizer was applied using the “CAUTAN” (Controlled Uptake Long Term Ammonium Nutrition) method. It was prepared by mixing one third ammonium sulfate and two thirds urea as nitrogen sources and by combining gypsum as a binder and loamy soil and compost as diffusion regulators in the beaker. In the first experiment, the application of NH_4 deposit fertilizer with 7.5 g gypsum as a binder resulted in increased tomato fruit yield and nitrogen uptake efficiency compared to control. In the second experiment, the application of NH_4 deposit fertilizer with loamy soil and compost as a diffusion regulator and adjusted C/N ratio to 16 also resulted in increased nitrogen uptake of fruits.

There are few growth studies evaluating within seasonal effects of N on vegetative growth and N accumulation of tomato conducted by Scholberg (2000). Growth analysis of field grown tomato for a number of Florida (USA) location and management systems are presented here. Severe N stress resulted in fewer and smaller, but thicker, leaves. With increasing N, average leaf area index increased from 0.75 to 3.0, but radiation use efficiency (RUE) typically increased less than 30%. Lower RUE under N limited condition reflected a decrease in N concentration of the most recently matured leaves from 40 mg g^{-1} to as little as 15 mg g^{-1} . Over the life of well fertilized crops, leaf N concentrations dropped from 55 to 65 mg g^{-1} during initial growth to 20 to 35 mg g^{-1} at final harvest.

Corresponding N concentrations for fruit and for stems were 30 to 35 mg g⁻¹ and 15 to 25 mg g⁻¹. Severe N stress affected leaf and stem N concentrations most drastically, whereas N in fruits was less variable.

Hoffland *et al.* (1999) conducted an experiment on tomato plants with varying N daily in exponentially increasing amounts to a nutrient solution at different rates. Leaves of plants grown at low N availability had a high leaf C:N ratio (21 g/g). The level of soluble carbohydrates correlated positively with susceptibility, independent of the growth method. It is therefore suggested that the effect of N availability on susceptibility can be explained by variation in levels of soluble carbohydrates which hence may play a role in the infection process.

The effects of low and high water vapor deficit regimes and electrical conductivities of 3.8 or 4.8 mS/cm on the growth and N uptake of 7 month old tomatoes in nutrient film technique (NFT) were investigated for 3 months by Bellert *et al.* (1998). Growth and N uptake were not modified by treatments. N accumulated in the aerial biomass in proportion to the dry matter. Total N concentration of the foliage was relatively constant and richer than that of vascular organs and fruits. A model is proposed to link total N concentration to dry matter accumulation.

In field trials on a red ferrallitic soil in northern Havana in 1994-95, tomato cv. Campbell 28 plants were fertilized with 0, 60, 120, 180 or 240 kg N/ha, starting 38 days after sowing was conducted by Adjanohoun *et al.* (1996). Although increasing rates of applied N had no effect on average fruit weight, they significantly increased fruit numbers, although application of 240 kg N /ha was excessive and significantly reduced yield compared with 120 or 180 kg N/ha (at the highest yield, obtained with 180 kg N/ha, was 38 t/ha). A mathematical expression describing the curve of yield response is presented, and from it the



optimum application rate was determined to be 158 kg N/ha, giving a fruit yield of 38.9 t/ha.

Tomato cv. Momotaro was grown using the nutrient film technique (NFT) in $\frac{1}{2}$ and $\frac{3}{4}$ to full strength Enshi shoho balanced feed (Hohjo *et al.* 1995). In the first experiment, nutrient solutions were adjusted to contain $\text{NO}_3^- \text{N} : \text{NH}_4^+ \text{N}$ ratios of 10:0, 9:1 and 8:2. Shoot and root (FW) were increased by an increasing proportion of $\text{NH}_4\text{-N}$ with both strengths of solution, whereas Ca and Mg uptake were decreased by an increasing proportion of $\text{NH}_4^+ \text{N}$ only with the higher solution strength. Total yield was reduced by increasing the proportion of $\text{NH}_4^+ \text{N}$, particularly with the higher strength of solution, a combination that also caused a marked increase in the incidence of blossom –end rot (BER). In the second experiment, $\text{NO}_3^- \text{N} : \text{NH}_4^+ \text{N}$ ratios of 10:0 and 8:2 and Ca concentrations of 2, 4 and 6 meq /litre were used. The higher proportion of $\text{NH}_4^+ \text{N}$ significantly increased shoot and root FW, incidence of BER and leaf contents of N, P and K, whilst decreasing the leaf content of Ca. Increasing the Ca content of the medium caused an increase in early yield and leaf Ca content, and decrease in BER and leaf Mg content. The combination of 8:2, $\text{NO}_3^- \text{N} : \text{NH}_4^+ \text{N}$ and the lowest Ca concentration reduced total yield and leaf Ca content and significantly increased blossom end rot (BER).

Trpevski *et al.*(1992) carried out in trials with 3 rates N was applied at 0, 40, 80 and 120 kg/ha to soil manured with 40 t FYM /ha in early spring. The 2 higher N rates increased the yield of San Pjer but reduced the yield of the other 2 cultivars. The effects of treatments on fruit N, dry matter, organic acid and vitamin C contents were generally not significant.

An experiment was conducted by Kooner and Randhaoa (1990) at Punjab Agricultural University, Ludhiana to study the interaction of rates on sources of N with cultivars on the yield and processing quality of tomatoes in winter and spring seasons. Four rates of N (50, 100, 150 and 200 kg/ha) were applied as 2 sources,

viz. calcium ammonium nitrate (CAN) and urea, in a randomized, split plot design. PC produced significantly higher yields (222.7 kg/ha) than PK (208.9kg/ha) in the spring planting while in the winter planting OS (163.09 kg/ha) and CS (113.9 kg/ha) were the best. Yields increased linearly with increasing N rate up to 150 kg/ha and calcium ammonium nitrate(CAN) was the best source of N. Total Soluble Solid (TSS), juice percentage, ascorbic acid content and titratable acidity increased with increasing N up to 150 kg/ha.

In a study, on the effect of nitrogen fertilization and plant intensification, Midan *et al.* (1998) observed that increasing nitrogen rates linearly increased the number of fruits per plant. However, medium and higher nitrogen rates gave best total yield. With different nitrogen rates, three times of application improved fruit per plant weight and total yield.

Patil and Bojappa (1984) conducted an experiment to study the effects of cultivars and graded levels of nitrogen and phosphorus on certain quality attributes of tomato. The experiment consisted of the cultivars Pusa Ruby, Sious and Sweet-72. The plant received nitrogen @ 70, 110 and 150 kg/ha and phosphorous (P) @ 44 or 61.6 kg/ha with basal dressing of potassium (K) @ 49.8 kg/ha and FYM @ 25 t/ha. The highest fruit content of total sugars and next highest dry matter content were in sweet-72, while juice percentage was highest in pusa Ruby. Rising nitrogen rates increased fruit total increased fruit total sugars and juice percentage but decreased the dry matter content. Phosphorous had no appreciable effects as any of the indices studied.

Belichki (1984) reported that nitrogen was the most important nutrient. Flower and fruit numbers per plant were increased by nitrogen up to 240 kg /ha and fruit size was greatest 120 kg/ha.

Staneve (1983) conducted an experiment to investigate the effect of nitrogen supply on photosynthesis, leaf area and total dry matter in tomato and found that photosynthesis was inhibited by N deficiency. Leaf development and dry matter accumulation were greatest @10 meq/L N and declined at higher concentrations.

2.2 Effect of Phosphorous on the growth and yield of tomato

A short term experiment was conducted by Kaya and Higgs (2001) with tomato cultivars Blizzard, Liberto and Calypso was carried out in a controlled temperature room to investigate the effectiveness of phosphorus (P) and iron (Fe) supplemented in nutrient solution on plant growth at high zinc concentration. Application of supplementary P and Fe resulted in marked increases in both dry weight and chlorophyll concentrations achieving values not significantly different to the control. Application of supplementary P and Fe decreased Zn concentration in the leaves and roots of plants grown at high Zn, but Zn concentrations were still at toxic levels. Phosphorus and Fe concentration in leaves declined to a deficient level in the high Zn treatment, but was markedly increased in the roots. Application of supplementary P and Fe corrected both P and Fe deficiencies in leaves of plants grown at high Zn and reduced root P and Fe concentrations.

Silva *et al.* (2001) carried out a field experiment was at Lavras, Minas Gerais, Brazil during 1997 to study nutrient absorption and yield by tomato plants pruned and grown under high planting density with various gypsum (0.3, 0.6, 0.9 and 1.2 t/ha) and P_2O_5 (0.2, 0.4, and 0.6t/ha). A linear increase was observed as gypsum rates increased. Leaf content decreased from 3.0 to 1.7 g/kg with an increase in gypsum rates and leaf Mg content averaged 4.7 g/kg.

Kaya *et al.* (2001) conducted an experiment with three tomato cultivars grown hydroponically in a controlled temperature room for 6 weeks to investigate the effectiveness of foliar application of supplementary potassium and phosphorus to leaves of plants grown at high NaCl concentration (60 mM). Supplementary 5 mM

K and P as KH_2PO_4 was supplied via leaves to the plants grown at high NaCl (60 mM) twice a week for 4 weeks. The plants grown at high NaCl produced less dry matter and chlorophyll than those at normal nutrient solution for all three cultivars. Membrane permeability increased with high NaCl application and these increases in membrane permeability were decreased by supplementary K and P. Concentration of P and K were at deficient ranges in the plants grown at high NaCl levels and these deficiencies were corrected by supplementary K and P application via leaves.

Field experiments conducted by Nanadal *et al.* (1998) from 1989 to 1991 in Haryana, India, four levels of each of phosphorus and potash with tomato showed that increasing levels of phosphorus up to 50 kg P_2O_5 and potash up to 80 kg $\text{K}_2\text{O}/\text{ha}$ improved the height of plant, number of flowers, weight of fruit, early and total yield, ascorbic acid content, total soluble solids(TSS) and reducing and non reducing sugars in the fruit. Though the higher doses of 75 kg $\text{P}_2\text{O}_5/\text{ha}$ and 120 kg $\text{K}_2\text{O}/\text{ha}$ increased fresh and dry weight of plant and advanced flowering and fruiting, they deteriorated the quality of fruit.

Song *et al.* (1998) carried out an experiment to find out the effects of phosphate starvation on growth and P uptakes of tomato seedlings were studied. The seedlings were arranged to grow in complete (P adequate) or P depleted nutrient solution. The results revealed that, under P starved conditions average height of the seedlings decreased, while the main roots of the seedling were longer than those of the controls, especially under phosphate starvation for 5 and 7 days. Therefore the root : shoot ratios of the seedlings under starved conditions were also higher than those of the controls. In addition the early days of phosphate starvation had no obvious effect on the fresh weight accumulation by the seedlings, while the late days of starvation did reduce fresh weight accumulation.

Bar Yosef (1995) conducted an experiment of nutrient uptake and dry matter production by a green house tomato (cv-144) crop was investigated in response to the concentration of P (0, 10, 30 and 60 mg P/litre) in the liquid feed and chemical properties. The leaf phosphorous content and dry matter production were increased significantly with increasing applied phosphorus. With increasing phosphorous concentration in leaves, there was an increase in leaf nitrogen and a decrease in potassium that become more pronounced above 0.65 g P/100g plant dry weight. With dry matter production, there was a marked fall-off below a leaf P concentration of 0.3 g P/100g plant dry weight.

Topeuoglu and Yalcin (1994) observed that different rates of lime (calcium hydroxide) and phosphorus were applied to potted tomato plants. Increasing rates of lime application increased individual fruit weight but decreased fruit yield/plant. Increasing rates of phosphorous increased vegetative growth and fruit yields.

Candilo *et al.* (1993) carried out an experiment to study the response of sulphur, phosphorus and application led to significant linear increases in cloth marketable yield and TSS content. In all interactions, the highest yields were obtained with 450 kg S, K₂O and P₂O₅/ha.

2.3 Integrated effect of nitrogen and phosphorous on the growth and yield of tomato

Garton and Widders (1990) carried out an experiment on seedlings of processing tomato cv. H 2653 grown in plug trays in a soil less growing medium. Application of nutrient solutions containing 10 or 20 mM N and 2 or 5 mM P for 10 days before transplanting altered the total ammoniacol N and P, and the soluble NO₃⁻ N and PO₄³⁻ P concentrations in the shoot tissue at transplanting. Post-transplanting shoot and root growth were more rapid in late May, plantings than in earlier plantings. The 20 mM N and 2mM P pre-transplant treatment caused the most



rapid shoot growth following early season planting in the field at Harrokw. Rapid seedling establishment after transplanting was generally not a good indicator of potential fruit yield. Withholding fertilizer temporarily before transplanting resulted in depletion in tissue N and P concentrations, slow post-transplanting shoot growth, and lower yields.

Tomato cv. Hisar Arun and okra cv. Pusa Sawani were planted during the 1991-92 and 1992-93 seasons at Haryana Agricultural University India, as a crop after potato by Taya *et al.* (1994). The preceding tomato crop received either the recommended fertilizer regime of 150 kg N/ha, 50 kg P_2O_5 /ha and 100 kg K_2O /ha (F_1). The subsequent crops received either N fertilizer @ 0 (N_0), 25 (N_1), 50 (N_2) of the recommended rate or 100 kg N/ha, 50 kg P_2O_5 /ha and 50 kg K_2O /ha (NPK). The F_1 treatments increased plant height @ 90 days after planting, fruit number and yield in the subsequent crops of tomato and okra; the F_1 treatment also increased the NPK content of foliage, and delayed flowering in both crops. With the tomato crop, there were no significant differences in fruit number and yield. Leaf N concentration increased with increasing N rate; N application had an effect on leaf P and K contents.

Field experiments were conducted on at Uttaranchal, India, by Singh *et al.* (2004) to determine the effects of integrated nutrient management on crop nutrient uptake and yield under okra-pea-tomato cropping sequence. In the sequence, treatments were given to okra crop, while in the succeeding crops (pea and tomato), only recommended dose of fertilizers were applied on the basis of soil test. The treatments consisted of NPK recommended dose of 80:30:30 kg/ha (T_1); farmyard manure (FYM) @15 tones/ha + rest of the NPK (T_2); neem cake @ 3 q/ha +rest of the NPK (T_3); poultry manure @ 3 tonnes/ha + rest of the NPK (T_4); *Azospirillum* +75% N +recommended dose of P and K (T_5); vesicular arbuscular mycorrhiza (VAM) + 50% P + recommended dose of N and K (T_6); phosphate solubilizing bacteria (PSB) +75% P + recommended dose of N and K (T_7); *Azospirillum*

+VAM+PSB+ rest of the NPK (T_8); micronutrient + recommended dose of NPK (T_9); FYM + *Azospirillum* +VAM+PSB+ rest of the NPK (T_{10}); and recommended dose of NPK + pea straw incorporation in the soil before tomato planting (T_{11}). In case of okra and pea crops, only the recommended dose of NPK was given in T_{11} . The treatments were applied in the first crop and their effect was observed on instant as well as succeeding crops. The integrated use of organic and inorganic sources of nutrients and bio-fertilizers increased the N, P and K concentrations in the plants of okra, pea and tomato. The integrated nutrient management also significantly increased shoot dry matter yield of tomato and fruit yields of okra and tomato.

A field experiment was undertaken in Peshawar, Pakistan in the summer of 1995-96 by Khalil *et al.* (2001) to determine the appropriate nitrogen fertilizer for maximum tomato yield and its effects on various agronomic characters of tomato. Treatments comprised: untreated control; 150 kg ammonium nitrate/ha; 150 kg ammonium nitrate/ha+100 kg P/ha+50 kg K/ha; 150 kg ammonium sulphate; 150 kg ammonium sulphate/ha+100 kg P/ha+50 kg K/ha; 150 kg urea/ha; 150 kg urea/ha+100 kg P/ha +50 kg K/ha. Generally, ammonium sulfate fertilizer was the most efficient source of nitrogen for tomato production followed by urea and ammonium nitrate. The ammonium sulfate +P +K treatment was the best among all treatments with respect to days to flower initiation (57 days), days to first picking (94 days), weight of individual fruit (50.8 g), weight of total fruits per plant (1990 g) and yield (21865 kg/ha). The control resulted in the significantly lowest response with respect to different agronomic characters under study.

Mohanty and Hossain (2001) conducted a field experiment during rabi 1994/95, 1995/1996 and 1996/97 at Bhawanipatna, Orissa, India to investigate the effect of nitrogen, potash and macronutrient application on the yield and yield attributes of tomato. Twelve different treatment combinations were tested and data were recorded for fruit yield, fruit weight, and benefit-cost ratio. Two treatments were found superior over the other treatments and gave the highest yields with high

benefit cost ratio over the 3 year study period; (i) 180 kg N/ha +60 kg P/ha+60 kg K/ha with top dressing of N twice at 30 and 60 days after planting (DAP); (II) 120 kg N/ha+60 kg P/ha +60 kg K/ha with top dressing of N twice at 30 and 60 DAP along with two sprays of macronutrients at flowering and 15 days thereafter.

Felipe and Casanova (2000) carried out an experiment to determine the effects of N (0, 90, 180 and 270 kg/ha), P (P_2O_5 , 0, 135, 270 and 405 kg/ha), and K (K_2O , 0, 90, 180 and 270 kg/ha) on the yield and number of tomato were investigated in the field in Venezuela. The best treatment, with the highest yield and number of fruits per plant, was 180 kg N, 270 kg P_2O_5 , and 180 kg K_2O /ha. It was possible to decrease the application of nutrients, particularly P. The increased yield was not due to larger fruits, but to an increased yield was not due to larger fruits, but to an increase in the number of fruits.

A field experiment was conducted by Sharma *et al.* (1999) with involving 4 levels of nitrogen (100, 150, 200 and 250 kg N/ha), 3 levels of phosphorus (60, 120 and 180 kg P_2O_5) and 3 hybrids (Naveen, MTH-16 and Rupali) and a local cultivar (Solan Gola) was conducted at Solan, India, to study the response of tomato hybrids to N and P. All the hybrids gave significantly higher total fruit yields than the local cultivar. Naveen recorded the greatest total fruit yield, while remaining statistically at par with MPH-16 and Rupali. Application of 200 kg N /ha resulted in significantly greater fruit size and mean fruit weight, weight, compared to the other application rates. A significant improvement in plant height, fruit size and total fruit yield was observed with the application of phosphorus from 60 to 180 kg P_2O_5 /ha.


A fertilizer trial tomato cultivars Acc-99 and Sweet-72 was carried out by Pandey *et al.* (1996) to study the effect of nitrogen and phosphorous at Jabalpur, India. Using nitrogen rate 0, 40, 80 or 120 kg /ha and phosphorous 0, 40 or 80 kg/ha, they concluded that the fruit yield was increased as nitrogen (N) rate increased up to 80 kg/ha. There was no significant difference between cultivators for fruit yield but there was an interaction nitrogen and phosphorus rates. Overall the highest

yield (49.95 t/ha) was obtained from Acc-90 given 80 kg/ha each of nitrogen and phosphorous.

Cerne and Briski (1993) conducted field trials on the fertilizer and irrigation requirement of tomato cv. Rudgers plants where 250 kg N and 725 kg P_2O_5 /ha plus 200 or 400 kg K_2O /ha in the first year, 0 or 200 kg K_2O /ha in the second year and 0 or 40 ton stable manure/ha were received all treatments. Plots were irrigated or non-irrigated (i.e. rainfed only). The combination of 400 kg K_2O /ha, stable manure and irrigation gave the highest total yield in the first and second years (1.03 and 2.25 kg/plant, respectively). The percentages of class 1 fruits were increased by irrigation in all cases.

The effects of adding Zn (5 kg/ha) Cu (3 kg/ha) or FYM (30 t/ha) to the basic N:P:K (222:160:100 kg/ha) treatment as leaf transpiration and chlorophyll content and fruit ascorbic acid and sugar contents were studied by Annanurova *et al.* (1992). The treatment was generally beneficial and the number and mean weight of fruits were increased. Application of NPK alone increased yield/plant 43.4%, compared with the untreated control.

While studying the nitrogen, phosphorous and potassium fertility regimes affect tomato transplant growth, Melton and Dufault (1991) found that as nitrogen increased from 25 to 225 mg/litre, fresh shoot weight, plant height, stem diameter leaf number, leaf area shoot and root dry weight and total chlorophyll content of seedlings ready for transplanting increased. Phosphorous @ 45 mg/litre increased fresh shoot weight, plant height, stem diameter, leaf number are compared with 5 to 15 mg P/litre. Potassium did not significantly influence any of the growth variables measured in the study.



Chapter III

Materials and Methods

CHAPTER III

MATERIALS AND METHODS

The experiment was conducted at the Research farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka, during November 2008 to March 2009 to find out the effect of nitrogen (N) and phosphorous (P) on the growth and yield of BARI Tomato-2 which is one of the high yielding varieties of indeterminate type.

3.1 Experimental site

The experiment was carried out in the experimental field of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka (Appendix I). The experimental site was located at $23^{\circ}77'$ N latitude and $90^{\circ}3'$ E longitude.

3.2 Characteristics of soil

The soil of the experimental site belongs to Tejgaon series under the Agro-ecological zone, Madhupur Tract (AEZ-28), which falls into Deep Red Brown Terrace Soil. The characteristics of the soil under the experimental plot were determined in the Soil Testing Laboratory (SRDI) Khamarbari, Dhaka and details of the recorded soil characteristics were presented in Table 1 and 2.

Table 1. Morphological characteristics of the experimental field

Morphological Features	Characteristics
Location	Sher-e-Bangla Agricultural University Farm, Dhaka-1207
AEZ	AEZ-28,(Modhupur Tract)
General Soil Type	Deep Brown Terrace Soil
Soil Series	Tejgaon
Topography	Fairly leveled
Depth of Inundation	Above flood level
Drainage condition	Well drained
Land Type	High land

Table 2. Physical and chemical properties of the experimental initial soil

Soil properties	Value
A. Physical properties	
1. Partical size analysis of soil	
Sand (%)	27.50
Silt (%)	40.30
Clay (%)	32.20
2. Soil texture	Silty-clay
3. Bulk density (g/cm^3)(0-30 cm depth)	1.39
4. Particle density (g/cm^3)(0-30 cm depth)	2.55
B. Chemical properties	
1. Soil pH	5.00
2. Organic carbon (%)	0.65
3. Organic matter (%)	1.23
4. Total Nitrogen (%)	0.109
5. C:N ratio	9.75:1
6. Available P (ppm)	0.24
7. Exchangeable K (me/100 g soil)	10.79

3.3 Climate

The experimental area has sub-tropical climate characterized by heavy rainfall during May to September and scant rainfall during rest of the year. The annual precipitation of the site is 2152 mm and potential evapotranspiration is 1297 mm. The average maximum temperature is 30.34⁰ C and average minimum temperature is 21.21⁰ C. The average mean temperature is 25.17⁰ C. The experiment was done during the rabi season. The temperature during the cropping period ranged from 12.2⁰ C to 29.20⁰ C. The humidity varied from 71.52% to 81.20%. The day length was reduced to 10.50 to 11.0 hours only and there was a very little rainfall from the beginning of the experiment to harvesting.

3.4 Seeds and variety

BARI Tomato-2(Raton), a high yielding variety of tomato (*Lycopersicon esculentum* Mill.) was used in this experiment developed by Bangladesh Agricultural Research Institute (BARI) Gazipur. The seeds were collected from Horticultural Research Centre (HRC) of BARI, Gazipur.

3.5 Raising of seedlings

The selected land for nursery bed was well drained and sandy loam type soil. The area was well prepared and converted into loose friable and dried mass to obtain fine tilth. All weeds and dead roots were removed and the soil was mixed with well rotted cow dung at the rate of 5 ton/ha. The size of the seed bed was 3m × 3m, raised 10 to 12 cm (approximately) above the ground level. The seed bed was prepared for raising the seedlings. Ten (10) grams of seeds were sown in the seed bed on 25 October, 2008. After sowing, the seeds were covered with light soil. Miral 3-GN was applied in seed bed as precautionary measures against ants and worms. Complete germination of the seeds took place with 5 days after seed sowing. Necessary shading was made by bamboo mat (chatai) from scorching sunshine or rain. No chemical fertilizer was used in the seed bed.

3.6 Treatments

The experiment was laid out in a Randomized Complete Block Design (RCBD) with three replications of each fertilizer treatment combinations. Fertilizer treatments consisting of 4 levels of N and P_2O_5 on the growth and yield of tomato. The study comprised the following treatments:

A. Four levels of nitrogen (N)

- i. N_0 : 0 kg N/ha
- ii. N_{100} : 100 kg N/ha
- iii. N_{150} : 150 kg N/ha
- iv. N_{200} : 200 kg N/ha

B. Four levels of phosphorous (P_2O_5)

- i. P_0 : 0 kg P_2O_5 /ha
- ii. P_{50} : 50 kg P_2O_5 /ha
- iii. P_{100} : 100 kg P_2O_5 /ha
- iv. P_{150} : 150 kg P_2O_5 /ha

There were 16 (4×4) treatment combinations. The treatment combinations were as follows

1. N_0P_0 = Control (without N and P application)
2. N_0P_{50} = 0 kg N /ha+ 50 kg P_2O_5 /ha
3. N_0P_{100} = 0 kg N /ha +100 kg P_2O_5 /ha
4. N_0P_{150} = 0 kg N /ha +150 kg P_2O_5 /ha
5. $N_{100}P_0$ = 100 kg N /ha + 0 kg P_2O_5 /ha
6. $N_{100}P_{50}$ =100 kg N /ha + 50 kg P_2O_5 /ha
7. $N_{100}P_{100}$ =100 kg N /ha + 100 kg P_2O_5 /ha
8. $N_{100}P_{150}$ =100 kg N /ha + 150 kg P_2O_5 /ha
9. $N_{150}P_0$ =150 kg N /ha + 0 kg P_2O_5 /ha
10. $N_{150}P_{50}$ =150 kg N /ha + 50 kg P_2O_5 /ha
11. $N_{150}P_{100}$ =150 kg N /ha + 100 kg P_2O_5 /ha
12. $N_{150}P_{150}$ = 150 kg N /ha + 150 kg P_2O_5 /ha
13. $N_{200}P_0$ =200 kg N /ha + 0 kg P_2O_5 /ha

$$14. N_{200}P_{50} = 200 \text{ kg N /ha} + 50 \text{ kg P}_2\text{O}_5/\text{ha}$$

$$15. N_{200}P_{100} = 200 \text{ kg N /ha} + 100 \text{ kg P}_2\text{O}_5/\text{ha}$$

$$16. N_{200}P_{150} = 200 \text{ kg N /ha} + 150 \text{ kg P}_2\text{O}_5/\text{ha}$$

Fertilizer treatments were randomly distributed in each block.

3.7 Design and layout of experiment



The two factors experiment was laid out in a Randomized Complete Block Design (RCBD) with three replications. An area of 335.7 m² (37.3 m × 9.0 m) was divided into three equal blocks. The layout of the experiment was prepared for distributing the treatment combinations into the every plot of each block. Each block was divided into 16 plots where 16 treatment combinations were allotted at random. There were 48 unit plots altogether in the experiment. Each of the plot area was 3.60 m² (2m × 1.8m). The distances maintained between two blocks and two plots were kept 1.0 m and 0.5 m, respectively. The layout of the experiment is shown in Figure 1.

3.8 Collection and preparation of initial soil sample

The initial soil samples were collected before land preparation from a 0-15 cm soil depth. The samples were drawn by means of an auger from different location covering the whole experimental plot and mixed thoroughly to make a composite sample. After collection of soil samples, the plant roots, leaves etc. were picked up and removed. Then the samples were air-dried and sieved through a 10-mesh sieve and stored in a clean plastic container for physical and chemical analysis.

3.9 Land preparation

The selected land for the experiment was first opened 25 November 2008 by a tractor. The land was well prepared with the tractor followed by harrowing and laddering up to a good tilth. All weeds and stubbles were removed. Making ridges (20 cm high) around each plot to restrict the lateral run off of irrigation water. The individual plots were finally prepared through addition of the basal doses of manure and fertilizers of that respective treatment.



Block to Block distance = 1m
 Plot to Plot distance = 50 cm
 Unit Plot size = $2\text{m} \times 1.8\text{m} = 3.6\text{m}^2$
 Plant to plant distance = 50cm
 Row to row distance = 60cm
 Total area = $37.3\text{m} \times 9\text{m} = 335.7\text{m}^2$

Variety:
 BARI-Tomato-2(Ratan)

Treatment:

Four levels of nitrogen (N)

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

Four levels of phosphorous (P₂O₅)

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Figure1: Field layout of the experimental plot



3.10 Application of manure and fertilizers

The sources of N, P₂O₅, K₂O, S as urea, triple super-phosphate (TSP), muriate of potash (MOP), and gypsum were applied, respectively. One third (1/3) of whole amount of urea, the entire amounts of TSP, MOP and gypsum were applied during the final land preparation. The remaining urea were top dressed in two equal installments at 20 days after transplanting (DAT) and 35 days after transplanting, respectively. Well rotten cowdung 10 t/ha also applied during final land preparation.

3.11 Transplanting of seedlings

55 (06) 09/08/11
38728
Healthy and uniform sized 30 days old seedlings were uprooted from the seed bed and were transplanted in the experimental field on 28 November, 2008 maintaining a spacing of 60 cm and 50 cm between the rows and plants, respectively. The seed bed was watered before uprooting the seedlings so as to minimize the damage of the roots. This operation was carried out during late hours in the evening. The seedlings were watered after transplanting. Shading was provided by piece of banana leaf sheath for three days to protect the seedlings from the direct sun light. A strip of the same crop was established around the experimental field as border crop to do gap filling and to check the border effect.

3.12 Irrigation and drainage

Over-head irrigation was provided with a watering cane to the plots once immediately after seedling transplanting in every alternate day in the evening up to establishment of seedling. Further irrigation was done when needed. Stagnant water was effectively drained out at the time of heavy rain.

3.13 Gap filling and weeding

When the seedlings were established, the soil around the base of each seedling was pulverized. A few gap filling were done by healthy plants from the border whenever it was required. Weeding was done to keep the plots free from weeds,

easy aeration of soil, which ultimately ensured better growth and development. The newly emerged weeds were uprooted carefully. Breaking the crust of the soil was done when needed.

3.14 Pest management

The crop was infested with cutworm, leaf hopper and others. The insects were controlled successfully by spraying Malathion 57 EC @ 2 ml/L water. The insecticide was sprayed fortnightly from one week after transplanting to one week before first harvesting. Miral 3-GN was also applied during final land preparation as soil amendments. During foggy weather precautionary measures against disease infestation especially late blight of tomato was taken by spraying Dithane M-45 fortnightly @ 2 g/L water.

3.15 Harvesting

Fruits were harvested at 5 days intervals during maturity to ripening stage. The maturity of the crop was determined on the basis of red coloring of fruits. Harvesting was started from 2 March, 2009 and completed by 22 March, 2009.

3.16 Collection of experimental data

Five (5) plants from each plot were selected randomly and were tagged for the data collection. The sample plants were uprooted and dried properly in the sun. Data were recorded on the following parameters from the sample plants during the course of experiment.

- 1) Plant height (cm) at 30 DAS and 60 DAS and at final harvest
- 2) Number of leaves per plant at 30 DAS and 60 DAS
- 3) Number of branches per plant at harvest
- 4) Number of cluster per plant
- 5) Number of flowers per cluster
- 6) Number of flowers per plant
- 7) Number of fruits per cluster

- 8) Number of fruits per plant
- 9) Fruit size (length $\times$ diameter)
- 10) Weight of individual fruit (gm)
- 11) Fruit yield per plant (gm)
- 12) Fruit yield per plot (kg)
- 13) Fruit yield per hectare (ton)
- 14) Nitrogen uptake by shoot
- 15) Phosphorous uptake by shoot
- 16) Nitrogen uptake by fruit
- 17) Phosphorous uptake by fruit
- 18) Nitrogen content in post harvest soil
- 19) Phosphorous content in post harvest soil

1. Plant height

Plant height at 30 DAS, 60 DAS and final harvest stages was measured respectively from sample plants in centimeter from the ground level to the tip of the longest stem and mean value was calculated.

2. Number of leaves per plant at 30 DAS and 60 DAS

Number of leaves per plant at 30 DAS and 60 DAS was counted from sample plants and their average was taken as their number of total leaves per plant.

3. Number of branches per plant at harvest

Number of main branches per plant at final harvest was recorded from sample plants and the average value was taken.

4. Number of cluster per plant

The number of clusters was counted from the sample plants and the average number of cluster produced per plant was recorded at the time of final harvest.

5. Number of flowers per plant

The number of flowers per plant was counted from the sample plant after end of flowering stage and the average number of flowers was taken.

6. Number of fruits per plant

The number of fruits per plant was counted from the sample plant after end of fruit setting and the average number of fruits was taken.

7. Number of flowers per cluster

It was calculated by the following formula

$$\text{Number of flowers} = \frac{\text{Total number of flowers of sample plants}}{\text{Total number of clusters of sample plants}}$$

8. Number of fruits per cluster

It was calculated by the following formula

$$\text{Number of fruits per cluster} = \frac{\text{Total number of flowers of sample plants}}{\text{Total number of clusters of sample plants}}$$

9. Rate of fruit setting (%)

It was calculated by the following formula

$$\text{Rate of fruit setting (\%)} = \frac{\text{Total number of fruits per plants}}{\text{Total number of flowers per plants}} \times 100$$

10. Fruit size (length × breadth)

The length of fruit was measured with a slide calipers from the neck of the fruit to the bottom of 5 randomly selected marketable fruits from each plot, at each harvest. Their average length was recorded. The breadth of fruit was measured at the middle position of 5 randomly selected marketable fruits from each plot at each harvest with a slide calipers.

11. Weight of individual fruit (gm)

Among the ten harvest of marketable fruits during the period from first to final harvest, the first and last two harvest were omitted and ten intermediate harvest were used for individual fruit weight in gram (gm) by the following formula

Weight of individual fruit (gm) =

$$\frac{\text{Total weight of marketable fruits from 10 harvested sample plant}}{\text{Total number of marketable fruits from 10 harvested of sample plant}}$$

12. Fruit yield per plant (gm)

It was calculated by the following formula

$$\text{Weight of fruits per plant (gm)} = \frac{\text{Total weight of fruits from 10 samples}}{10}$$

13. Fruit yield per plot (kg)

A pan scale balance was used to take the weight of fruits during the period from first to final harvest.

14. Fruit yield per hectare (ton)

It was calculated by the following formula

$$\text{Fruit yield per hectare (ton)} = \frac{\text{Fruit yield per plot (kg)} \times 10000}{\text{Area of plot in square meter} \times 1000}$$

3.17 Analytical methods of soil and plant sample analysis

The analytical methods of physico-chemical properties of soil, nutrient content and quality of tomato are shown in the table 3

3.18 Chemical analysis of plant samples

1. Preparation of plant samples

Ten selected plants per plot were collected immediately after harvest. The selected plant was threshed. Both plant and fruit were cleaned and dried in an oven 65°C for 48 hours. The dried samples were put into small paper bags and kept into desiccators till being used.

2. Digestion of plant samples with sulphuric acid

For N determination, an amount of 0.1g plant sample (plant was taken into a 100 ml Kjeldhal flask). An amount of 1.1 g catalyst mixture (K₂SO₄: CuSO₄. 5H₂O: Se =100:10:1), 2 ml 30% H₂O₂ and 3 ml conc. H₂SO₄ were added into the flask. The flask was swirled and allowed to stand for about 10 minutes followed by heating at

200°C. Heating was continued until the digest was clear, and colorless. After cooling, the contents were taken into a 100 ml volumetric flask and the volume was made with distilled water. A blank digestion was prepared in a similar way. This digest was used for determining the nitrogen contents of plants samples.

3. Digestion of plant with nitric per chloric acid mixture

An amount of 0.5 g of plant was taken into a dry clean 100 ml Kjeldahl flask, 10 ml of di-acid mixture (HNO_3 and HClO_4 at the ratio of 2:1) was added and kept for few minutes. Then, the flask was heated at a temperature rising slowly up to 200°C. Heating was instantly stopped as soon as the dense white fumes of HClO_4 occurred and after cooling; 6 ml of 6 N HCl were added to it. The content of the flask was boiled until they become clear and colorless. This digest was used for determining P, K, S and Zn.

4. Determination of element in the digest (Table 3)

Nitrogen, phosphorous, potassium and sulphur content in the digest were determined by similar method as described in soil analysis.

3.19 Soil Sample Analysis

1. Total nitrogen

Total nitrogen of soil was determined by Micro Kjeldahl method where soil was digested with 30% H_2O_2 , H_2SO_4 and catalyst mixture (K_2SO_4 : $\text{CuSO}_4 \cdot 5\text{H}_2\text{O}$: Se powder in the ratio of 100:10:1). Nitrogen in the digest was estimated by distillation with 40% NaOH followed by titration of the distillate trapped in H_3BO_3 with 0.01 N H_2SO_4 (Bremner and Mulvaney, 1982).



Table 3. Different methods for determination of physico-chemical properties of soil.

Physical properties of soil

Soil properties	Determination method
Soil texture (particle size analysis)	Hydrometer method
Bulk density	Core sampling method
Particle density	Pycnometer method
% Porosity	Relation between bulk density and particle density

Chemical properties of soil

Soil properties	Determination Method
Soil pH	1:2.5 soil water ratio, using glass electrode method.
Organic carbon	Wet oxidation method as outlined by Piper (1942): Through oxidation of organic matter with potassium dichromate and titrated with ammonium ferrous sulphate, using phenyl anthranolic acid indicator.
Total N	Micro-Kjeldahl method following H_2SO_4 digestion and steam distillation with NaOH. Ammonia thus collect in boric acid was determined titrimetrically.
Available P	Extracted by 0.5 M $NaHCO_3$ (pH 8.5) and determined colorimetrically using molybdate blue method (Olsen and Dean, 1965).
Exchangeable K	Extracted by ammonium acetate solution determined by atomic absorption spectrophotometer (Christian and Feldman, 1970).

2. Available phosphorous

Available phosphorous was extracted from soil by shaking with 0.5 M NaHCO_3 solution of pH 8.5 (Olsen *et al.*, 1954). The phosphorous in the extract was then determined by developing blue color using SnCl_2 reduction of phosphorous molybdate complex. The absorbance of the molybdophosphate blue color was measured at 660 nm wave length by spectrophotometer and available P was calculated with the help of standard curve.

3.20 Statistical Analysis

The data obtained for different parameters were statistically analyzed to find out the significance difference of the different levels of nitrogen and phosphorous on yield and yield contributing characters of tomato and nutrient uptake and content by plant and fruit. The mean values of all the characters were calculated and analysis of variance was performing by the 'F' (variance ratio) test. The significance of the difference among the treatment combinations means was estimated by the Duncan's Multiple Range Test (DMRT) at 5% level of probability (Gomez and Gomez, 1984).



Chapter IV

Results and Discussion



CHAPTER IV

RESULTS AND DISCUSSION

The experiment was conducted at the Research farm of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka, during November 2008 to March 2009 to find out the effect of nitrogen (N) and phosphorous (P) on the growth and yield of BARI Tomato-2 which is one of the high yielding varieties of indeterminate type. The experiment was carried out in the experimental field of Sher-e-Bangla Agricultural University, Sher-e-Bangla Nagar, Dhaka.

To determine the effect of different levels of nitrogen and phosphorous on yield and yield contributing characters of tomato as well as the nutrient content and their uptake by plants and fruits. Data on different characters were recorded and analyzed to find out the effects of nitrogen and phosphorous. The analyses of variance (ANOVA) of the data on different components are given in Appendix (II-VII). The results have been presented and discussed and possible explanations have been given under the following parameters:

4.1 Yield and yield contributing character of tomato

Yield contributing character such as plant height, number of leaves, number of branches, number of cluster per plant, number of flowers per cluster, number of flowers per plant, number of fruits per cluster, number of fruits per plant, rate of fruit setting (%), fruit size, weight of individual fruit, fruit yield per plant, fruit yield per hectare, yield increased over control, shoot and root dry weight etc. were recorded.

4.1.1 Plant height

Plant height varied significantly due to application of nitrogen in tomato under the present trial (Appendix III). At 30 days after transplanting (30 DAT) the maximum plant height (60.38 cm) was recorded from N₂₀₀ treatment consisting of

200 kg N/ha which was statistically identical (58.76 cm) with N_{150} treatment as 150kg N/ha and the minimum plant height (56.16 cm) was recorded from N_0 treatment that is control condition which was closely (57.25cm) followed by N_{100} treatment as 100kgN /ha (Table 4).

At 60 days after transplanting (60 DAT) the highest plant height (78.31cm) was recorded from N_{200} treatment consisting of 200 kg N/ha which was statistically identical (77.05cm) with N_{150} treatment as 150 kg N/ha and the lowest plant height (70.78 cm) was recorded from N_0 treatment (Table 4).

During the harvesting period, the maximum plant height (89.58 cm), was recorded from N_{200} treatment consisting of 200 kg N/ha and the minimum plant height (56.16cm) was recorded from N_0 treatment which was closely (80.17 cm) followed by N_0 treatment (Table 4).

Probably, 200 kg N/ha ensured the favorable condition for growth of tomato plant and the ultimate result is the tallest plant where as above this level of nitrogen hinder the growth and plant height decreases. Melton and Dufault (1991) found that plant height of tomato was increased as highest level of nitrogen. Similar result was reported by Chung *et al.* (1992), Grela *et al.* (1988), Sharma and Mann (1971)

Different level of phosphorous showed statistically significant differences for plant height (Appendix III). At 30 days after transplanting (30 DAT) the maximum plant height (59.43 cm), (Figure 3) was recorded from P_{100} treatment consisting of 100kg P_2O_5 /ha which was statistically identical (59.15cm) with P_{150} treatment as 150kg P_2O_5 /ha and (58.51cm) with P_{50} treatment as 50kg P_2O_5 /ha the minimum plant height (55.46 cm) was recorded from P_0 treatment that is control condition (Table .4).



At 60 days after transplanting (60 DAT) the maximum plant height (77.61 cm), was recorded from P_{150} treatment consisting of 150kg P_2O_5 /ha which was statistically identical(77.24 cm) with P_{100} treatment as 100kg P_2O_5 /ha and (58.51cm) with P_{50} treatment as 50kg P_2O_5 /ha the minimum plant height (72.046 cm) was recorded from P_0 treatment (Table 4). During the harvesting period, the maximum plant height (88.93 cm), was recorded from P_{150} treatment consisting of 150kg P_2O_5 /ha which was statistically identical (88.08) with P_{100} treatment as 100kg P_2O_5 /ha and the minimum plant height (80.64 cm) was recorded from N_0 treatment (Table 4).

Interaction effect was also recorded between nitrogen and phosphorous in consideration of plant height under the present experiment and found statistically significant variation (Appendix III). At 30 day after transplanting, (30 DAT) the maximum plant height (61.74 cm), was recorded from $N_{200} P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha which was statistically identical (60.94 cm) with $N_{200}P_{50}$ treatment as 200 kg N/ha +50 kg P_2O_5 /ha and (60.91 cm) with $N_{200}P_{100}$ treatment as 200 kg N/ha+ 100 kg P_2O_5 /ha. The minimum plant height (52.77 cm) was recorded from N_0P_0 treatment which was statistically identical (54.43 cm) by $N_{100}P_0$ treatment as 100 kg N /ha+0 kg P_2O_5 /ha (Table 5).

At 60 days after transplanting (60 DAT) the maximum plant height (81.48 cm), was recorded from $N_{200} P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha which was statistically identical (79.99 cm) (Figure 4) with $N_{200}P_{100}$ treatment as 200 kg N/ha +100 kg P_2O_5 /ha. The minimum plant height (68.23 cm) was recorded from N_0P_0 treatment which was statistically similar (70.51 cm) followed by N_0P_{50} treatment as 0 kg N /ha+50 kg P_2O_5 /ha (Table 5).



Table 4. Effect of nitrogen and phosphorous on plant height, number of leaves per plant and branches per plant of tomato

Treatments	Plant height (cm)			No. of leaves / plant at 60 DAT	No. of branches / plant at harvesting period
	30 DAT	60 DAT	Harvesting period		
Nitrogen					
N ₀	56.16c	70.78c	80.17d	39.72d	9.668c
N ₁₀₀	57.25bc	75.77b	85.05c	43.08c	10.44b
N ₁₅₀	58.76ab	77.05ab	87.47b	46.92b	10.90a
N ₂₀₀	60.38a	78.31a	89.58a	51.72a	11.26a
LSD	1.636	1.765	1.747	2.073	0.3964
Phosphorous					
P ₀	55.46b	72.04c	80.64c	40.10c	9.458c
P ₅₀	58.51a	75.00b	84.63b	45.08b	10.29b
P ₁₀₀	59.43a	77.24a	88.08a	47.78a	11.18a
P ₁₅₀	59.15a	77.61a	88.93a	48.49a	11.35a
LSD	1.889	2.038	2.017	2.393	0.4577

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability.

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Table 5. Interaction effect of nitrogen and phosphorous on plant height, number of leaves per plant and branches per plant of tomato

Treatments (N×P)	Plant height (cm)			No. of leaves/ plant at 60 DAT	No. of branches / plant at harvesting period
	30 DAT	60 DAT	Harvesting period		
N ₀ P ₀	52.77d	68.23h	75.44i	35.87e	8.39h
N ₀ P ₅₀	56.50bc	70.51gh	79.50h	39.52de	9.40g
N ₀ P ₁₀₀	58.55ab	72.85efg	82.80gh	41.50cd	10.32def
N ₀ P ₁₅₀	56.82bc	71.51fgh	82.95fgh	42.01cd	10.56def
N ₁₀₀ P ₀	54.43cd	72.38fg	80.40h	39.53de	9.69fg
N ₁₀₀ P ₅₀	58.15ab	75.35cdef	84.30efg	43.38cd	10.32def
N ₁₀₀ P ₁₀₀	58.25ab	77.38bcd	87.51cde	44.42c	10.77cde
N ₁₀₀ P ₁₅₀	58.17ab	77.97abcd	87.98cde	45.00c	10.99bcd
N ₁₅₀ P ₀	56.71bc	73.23efg	82.20gh	41.67cd	9.82fg
N ₁₅₀ P ₅₀	58.46ab	76.72bcde	86.71def	45.31c	10.57def
N ₁₅₀ P ₁₀₀	60.03ab	78.76abc	89.82bcd	49.84b	11.57abc
N ₁₅₀ P ₁₅₀	59.86ab	79.49ab	91.15abc	50.87b	11.66ab
N ₂₀₀ P ₀	57.92abc	74.34defg	84.52efg	43.31cd	9.93efg
N ₂₀₀ P ₅₀	60.94a	77.43bcd	88.00cde	52.10ab	10.87bcd
N ₂₀₀ P ₁₀₀	60.91a	79.99ab	92.20ab	55.36a	12.05a
N ₂₀₀ P ₁₅₀	61.74a	81.84a	93.62a	56.09a	12.19a
LSD	3.272	3.529	3.494	4.145	0.7927
CV (%)	3.37	2.80	2.45	5.48	4.50

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability.

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

During the harvesting period, the maximum plant height (93.62 cm), was recorded from $N_{200} P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha which was statistically identical (92.20 cm) with $N_{200}P_{100}$ treatment as 200 kg N/ha +100 kg P_2O_5 /ha. The minimum plant height (75.44 cm) was recorded from N_0P_0 treatment that is control condition which was similar to (79.50 cm) N_0P_{50} treatment as 0 kg N /ha+50 kg P_2O_5 /ha. (Table 5)

4.1.2 Number of leaves per plant at 60 DAT

Number of leaves per plant varied significantly due to application of nitrogen in tomato under the present trial (Appendix III). At 60 days after transplanting (60 DAT) the highest number of plant leaves (51.72), was recorded from N_{200} treatment consisting of 200 kg N/ha and the lowest number plant leaves (39.72) was recorded from N_0 treatment (Table 4) .The ranges of number of leaves per plant at 60 DAT incase of nitrogen application was 39.72 to 51.72

Different level of phosphorous showed statistically significant differences for plant leaves (Appendix III). At 60 days after transplanting (60 DAT) the number plant leaves (48.49), was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical with P_{100} treatment (47.78) as 100 kg P_2O_5 /ha and the minimum number of plant leaves (40.10) was recorded from P_0 treatment (Table 4). The ranges of number of leaves per plant at 60 DAT incase of phosphorous application was 40.10 to 48.49.

In the interaction effect, data recorded between nitrogen and phosphorous in consideration of plant leaves under the present experiment and found statistically significant variation (Appendix III). At 60 days after transplanting (60 DAT) the maximum number of plant leaves (56.09) was recorded from $N_{200} P_{150}$ treatment consisting of 200kg N/ha +150kg P_2O_5 /ha which was statistically identical(55.36) with $N_{200}P_{100}$ treatment as 200 kg N/ha +100 kg P_2O_5 /ha. The minimum number of



plant leaves (35.87) was observed from N_0P_0 treatment which was statistically similar (39.52) to the N_0P_{50} treatments as 0kgN /ha+50 kg P_2O_5 /ha (Table 5).

4.1.3 Number of branches per plant at final harvest

In response of number of branches per plant varied significantly due to application of nitrogen in tomato under the present trial (Appendix III). At final harvesting period, the large number of branches per plant (11.26), was observed from N_{200} treatment consisting of 200kg N/ha and which was statistically identical (10.90) with N_{150} treatment as 150 kg N/ha and the lowest number of branches per plant (9.668) was observed from N_0 treatment (Table 4). The ranges of number of branches per plant at final harvest incase of nitrogen application was 9.66 to 11.26.

Different level of phosphorous showed statistically significant differences for plant leaves (Appendix III). At final harvesting period, the large number of branches per plant (11.35), was recorded from P_{150} treatment consisting of 150kg P_2O_5 /ha which was statistically identical(11.18) with P_{100} treatment as 100kg P_2O_5 /ha and the minimum branches per plant (9.45) was recorded from P_0 treatment (Table .4). The ranges of number of branches per plant at final harvest incase of phosphorous application was 9.45 to 11.35.

Due to interaction effect, the recorded data between nitrogen and phosphorous in consideration of number of branches per plant found statistically significant (Appendix III). At final harvesting period the maximum number of branches per plant (12.19) was recorded from $N_{200} P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha which was statistically identical (12.05) with $N_{200}P_{100}$ treatment as 200 kg N/ha +100 kg P_2O_5 /ha. The minimum number of branches per plant (8.39) was recorded from N_0P_0 treatment which was statistically similar (9.40) followed by N_0P_{50} treatment as 0 kg N /ha+50 kg P_2O_5 /ha (Table 5).

Table 6. Effect of nitrogen and phosphorous on number of cluster / plant, number of flowers/ cluster, number of flowers/ plant, number of fruits/ cluster and number of fruits/ plant of tomato

Treatments	Number of flower cluster / plant	Number of flowers/ cluster	Number of flowers/ plant	Number of fruits/ cluster	Number of fruits/ plant
Nitrogen					
N ₀	6.35d	4.520b	30.41c	3.325b	21.75c
N ₁₀₀	7.537c	4.813a	37.46b	3.697a	27.86b
N ₁₅₀	8.432b	4.932a	42.41a	3.820a	32.33a
N ₂₀₀	8.800a	4.943a	44.63a	3.828a	33.77a
LSD	0.3142	0.1625	1.991	0.2059	1.687
Phosphorous					
P ₀	6.365c	4.452c4.	30.01c	2.525b	22.59c
P ₅₀	7.645b	4.805b	37.88b	3.630ab	27.89b
P ₁₀₀	8.530a	4.947ab	43.38a	3.787a	32.49a
P ₁₅₀	8.580a	5.003a	44.02a	3.727ab	32.75a
LSD	0.3628	0.1877	2.299	0.23783.	1.948

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Table 7. Interaction effect of nitrogen and phosphorous on number of flower cluster/ plant, number of flowers/ cluster, number of flowers/ plant, number of fruits/ cluster and number of fruits/ plant of tomato

Treatments (N×P)	Number of flower cluster / plant	Number of flowers/ cluster	Number of flowers/ plant	Number of fruits/ cluster	Number of fruits/ plant
N ₀ P ₀	5.01g	4.10e	23.52h	3.21d	16.08f
N ₀ P ₅₀	6.41f	4.52d	29.97fg	3.33bcd	21.35e
N ₀ P ₁₀₀	6.98def	4.69bcd	33.74def	3.52abcd	24.57de
N ₀ P ₁₅₀	7.00def	4.77abcd	34.39de	3.24cd	24.99de
N ₁₀₀ P ₀	6.32f	4.42de	28.89g	3.52abcd	22.21e
N ₁₀₀ P ₅₀	7.53cd	4.78abcd	36.99cd	3.68abc	27.71cd
N ₁₀₀ P ₁₀₀	8.00bc	4.91abc	40.28bc	3.75ab	30.00bc
N ₁₀₀ P ₁₅₀	8.30b	5.14a	43.66b	3.84a	31.54b
N ₁₅₀ P ₀	6.75ef	4.62cd	32.19efg	3.67abcd	24.77de
N ₁₅₀ P ₅₀	8.00bc	4.91abc	40.28bc	3.77ab	30.16bc
N ₁₅₀ P ₁₀₀	9.54a	5.10a	49.65a	3.89a	37.11a
N ₁₅₀ P ₁₅₀	9.44a	5.10a	49.14a	3.95a	37.29a
N ₂₀₀ P ₀	7.38cde	4.67bcd	35.46de	3.70abc	27.31cd
N ₂₀₀ P ₅₀	8.64b	5.01ab	44.29b	3.74ab	32.32b
N ₂₀₀ P ₁₀₀	9.60a	5.09a	49.86a	3.99a	38.30a
N ₂₀₀ P ₁₅₀	9.58a	5.0ab	48.90a	3.88a	37.17a
LSD	0.6284	0.3251	3.983	0.4118	3.375
CV (%)	4.84	4.08	6.15	6.74	7.00

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability.

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

4.1.4 Number of flower clusters per plant

Application of nitrogen in tomato under the present trial data were found statistically significant in respect of number of flower clusters per plant (Appendix IV). The no of largest number flower clusters per plant (8.800), was found from N_{200} treatment consisting of 200kg N/ha and on the other hand the lowest number of flower clusters per plant (6.350) was recorded from N_0 treatment. Found the similar result that is application of 240 kg N/ha ensured the favorable condition for growth of tomato plant and the ultimate result is the number of flower cluster per plant. Grela *et al.* (1988) put forward almost similar opinion (Table 6).

Different level of phosphorous data showed statistically significant differences for flower clusters per plant (Appendix IV). The maximum number of flower clusters per plant (8.580) was found from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical (8.530) with P_{100} treatment as 100 kg P_2O_5 /ha whereas the minimum number of flower clusters per plant (6.365) was recorded from P_0 treatment under the present trial. With increasing level of phosphorous, plant growth increases and the number of flower clusters per plant also increases (Table 6).

In the interaction effect, data was also recorded between nitrogen and phosphorous in consideration of plant flower clusters under the present experiment and found statistically significant variation (Appendix IV). The maximum flower clusters per plant (9.60) was recorded from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha +100 kg P_2O_5 /ha which was statistically identical (9.580) with $N_{200}P_{150}$ treatment as 200 kg N/ha +150 kg P_2O_5 /ha, (9.540) with $N_{150}P_{100}$ treatment as 150 kg N/ha +100 kg P_2O_5 /ha and (9.440) with $N_{150}P_{150}$ treatment as 150 kg N/ha +150 kg P_2O_5 /ha. The minimum flower clusters per plant (5.01) was recorded from N_0P_0 treatment that is control condition (Table 7).



4.1.5 Number of flowers per cluster

Number of flowers per cluster varied significantly due to application of nitrogen in tomato under the present trial (Appendix IV). The highest number of flowers per cluster (4.943) was observed from N_{200} treatment consisting of 200 kg N/ha, which was statistically identical with N_{150} treatment (4.932) consisting of 150 kg N/ha, (4.813) with N_{100} treatment consisting of 100 kg N/ha and the lowest number of flowers per cluster (4.520) was found from N_0 treatment (Table 6).

Different level of phosphorous application showed statistically significant difference for flowers no. per clusters (Appendix IV). The maximum number of flowers per clusters (5.003) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical with P_{100} treatment (4.947) as 100 kg P_2O_5 /ha and the minimum number of flowers per cluster (4.452) was recorded from P_0 treatment (Table 6).

Due to interaction effect, data was also recorded between nitrogen and phosphorous in consideration of plant flowers no. per cluster and found statistically significant variation (Appendix IV). The maximum number of flowers per cluster (5.140) was found from $N_{100}P_{150}$ treatment consisting of 100 kg N/ha +150 kg P_2O_5 /ha which was statistically identical with both $N_{150}P_{100}$ and $N_{150}P_{150}$ treatment (5.100) as 150 kg N/ha +100 kg P_2O_5 /ha, and as 150 kg N/ha +150 kg P_2O_5 /ha, respectively and also (5.09) with $N_{200}P_{100}$ treatment as 200 kg N/ha +100 kg P_2O_5 /ha. The minimum number of flowers per cluster (4.1) were recorded from N_0P_0 treatment which was statistically identical with the treatment $N_{100}P_0$ (4.42) as 100 kg N/ha +0 kg P_2O_5 /ha (Table 7).

4.1.6 Number of flowers per plant

Application of different doses of nitrogen fertilizer in tomato, number of flowers per plant varied significantly (Appendix IV). The maximum number of flowers per plant (44.63) was recorded from N_{200} treatment consisting of 200 kg N/ha, which

was statistically identical (42.81) with N_{150} treatment consisting of 150 kg N/ha, and the minimum number of flowers per plant (30.41) was recorded from N_0 treatment (Table 6). The ranges of number of flowers per plant in case of nitrogen application was 30.41 to 44.63.

Due to different level of phosphorous data showed statistically significant difference for flowers no. per plant (Appendix IV). The highest number of flowers per plant (44.02), was observed from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical with P_{100} treatment (43.38) as 100 kg P_2O_5 /ha and the lowest number of flowers per plant was recorded from P_0 treatment (30.01) (Table 6). The ranges of number of flowers per plant in case of phosphorous application was 30.01 to 44.02.

In respect of interaction effect of data was recorded between nitrogen and phosphorous in consideration of flowers number per plant and observed statistically significant variation (Appendix IV). The maximum number of flowers per plant (49.86) was recorded from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha + 100 kg P_2O_5 /ha which was statistically identical, with $N_{150}P_{100}$ (49.65), $N_{150}P_{150}$ (49.14) and $N_{200}P_{150}$ (48.90) treatment as 150 kg N/ha + 100 kg P_2O_5 /ha, 150 kg N/ha + 150 kg P_2O_5 /ha, and as 200 kg N/ha + 150 kg P_2O_5 /ha respectively. The minimum number of flowers per plant were recorded from N_0P_0 (23.52) treatment (Table 7).

4.1.7 Number of fruits per cluster

Number of fruits per cluster varied significantly due to application different doses of nitrogen in tomato under the present trial (Appendix IV). First it is important to mention that the number of fruits per cluster increased with increasing level of N. The highest number of fruits per cluster (3.828) was recorded from N_{200} treatment consisting of 200 kg N/ha, which was statistically identical with N_{150} treatment (3.820) consisting of 150 kg N/ha, (3.697) with N_{100} treatment consisting of 100

kg N/ha and the lowest number of fruits no cluster (3.325) was recorded from N_0 treatment (Table 6). The ranges of number of fruits per cluster incase of nitrogen application was 3.325 to 3.828.

Application of different levels of phosphorous showed statistically significant difference for fruits number per clusters (Appendix IV). The maximum number of fruits per clusters (3.787), was recorded from P_{100} treatment consisting of 100 kg P_2O_5 /ha which was statistically identical with P_{150} treatment (3.727) as 150 kg P_2O_5 /ha and with P_{50} treatment (3.630) as 50 kg P_2O_5 /ha. The minimum numbers of fruits per cluster (3.525) were recorded from P_0 treatment (Table 6). The ranges of number of fruits per cluster incase of phosphorous application was 3.525 to 3.787.

In this trial, interaction effect was also recorded between nitrogen and phosphorous in consideration of fruits number per cluster and found statistically significant variation (Appendix IV). The maximum number of fruits per clusters (3.990), was recorded from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha +100 kg P_2O_5 /ha which was statistically identical with $N_{150}P_{150}$ (3.950), $N_{150}P_{100}$ (3.890) and $N_{200}P_{150}$ (3.880) treatment as 150 kg N/ha +150 kg P_2O_5 /ha, as 150 kg N/ha +100 kg P_2O_5 /ha and as 200 kg N/ha +150kg P_2O_5 /ha, respectively. The minimum numbers of fruits per cluster were recorded from N_0P_0 treatment (3.21) (Table 7).

4.1.8 Number of fruits per plant

Number of fruits per plant varied significantly due to application of nitrogen in tomato (Appendix IV). The highest amount of fruits per plant (33.77) was recorded from N_{200} treatment consisting of 200 kg N/ha, which was statistically identical with N_{150} treatment (32.33) consisting of 150 kg N/ha, and lowest amount of fruits per plant was recorded from N_0 treatment (21.75) (Table 6). Nassar (1988) found that increased nitrogen level was tended to increase average number of fruits cluster per plant.

Due to different level of phosphorous showed statistically significant difference for fruits number per plant (Appendix IV). The maximum fruits number per plant (32.75) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical with P_{100} treatment (32.49) as 100 kg P_2O_5 /ha and the minimum fruits number per plant was recorded from P_0 treatment (22.59) (Table 6). With increasing levels of phosphorous, plant growth increases and which ensure maximum number of flowers and fruits per cluster as well as the number of fruits per plant.

Interaction effect was also recorded between nitrogen and phosphorous in consideration of fruits number per plant and found statistically significant variation (Appendix IV). The maximum number of fruits per plant was recorded from $N_{200}P_{100}$ treatment (38.30), consisting of 200 kg N/ha +100 kg P_2O_5 /ha, which was statistically identical with $N_{150}P_{150}$ (37.29), $N_{200}P_{150}$ (37.17) and $N_{150}P_{100}$ (37.11) treatments. The minimum numbers of fruits per plant (16.08) were recorded from N_0P_0 treatment (Table 7). These result revealed that combined higher dose of nitrogen and phosphorous is required for attaining better growth and the ultimate results was desirable growth with maximum number of flower and fruit per cluster and highest number of fruits per plant. Winsor *et al.*, (1962) reported that increased application of nitrogen and phosphorous has the positive effect on number of fruits per plant.

4.1.9 Weight of individual fruit

Weight of individual fruit per plant varied significantly due to application of nitrogen in tomato under the present trial (Appendix V). On the one hand the maximum weight of individual fruit per plant (83.77 gm) was recorded from N_{200} treatment consisting of 200kg N/ha, which was statistically identical (83.27 gm) with N_{150} treatment consisting of 150 kg N/ha, and on the other hand the minimum weight of individual fruit per plant (66.22 gm) was recorded from N_0 treatment that is zero level of nitrogen (Table 8). Incase of N the maximum fruit weight

(79.25 gm) was found at 200 kg N/ha which is agreement with Ahmed and Choudhury (1990).

Different level of phosphorous showed statistically significant differences for weight of individual fruit per plant in tomato (Appendix V). The maximum weight of individual fruit per plant (81.67 gm), was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical (80.96 gm) with P_{100} treatment as 100 kg P_2O_5 /ha and (79.49 gm) with P_{50} treatment as 50 kg P_2O_5 /ha and the minimum weight of individual fruit per plant (71.54) was recorded from P_0 treatment that is zero level of phosphorous (Table 7). The ranges of weight of individual fruit incase of phosphorous application was 71.54 gm to 81.67 gm

Incase of interaction effect, data was also recorded between nitrogen and phosphorous in consideration of weight of individual fruit per plant and found statistically significant variation (Appendix V). The maximum weight of individual fruit per plant (87.59 gm) was recorded from $N_{150}P_{100}$ treatment consisting of 150 kg N/ha +100 kg P_2O_5 /ha which was statistically similar with $N_{200}P_{150}$ (87.29 gm), $N_{150}P_{150}$ (87.29 gm) and $N_{200}P_{100}$ (86.97 gm) treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha and 150 kg N/ha +150 kg P_2O_5 /ha and (86.97 gm) from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha +100 kg P_2O_5 /ha, and the minimum weight of individual fruit per plant (63.85 gm) was recorded from N_0P_0 treatment (Table 8).

4.1.10 Fruit size (Fruit length)

Length of fruit incase of fruit size of tomato plant varied significantly due to application of nitrogen in tomato (Appendix V). The highest fruit length (5.320 cm) was recorded from N_{150} treatment consisting of 150 kg N/ha, which was statistically identical (5.278 cm) with N_{200} treatment consisting of 200kg N/ha, and (5.225 cm) with N_{100} treatment consisting of 100 kg N/ha, the minimum fruit



length (4.605 cm) was recorded from N_0 treatment that zero level of nitrogen (Table 8).

Due to different level of phosphorous, data showed statistically significant difference for fruit length of plant (Appendix V). The maximum fruit length (5.262 cm) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha and the minimum fruit length (4.960 cm) was recorded from P_0 treatment that is zero level of phosphorous (Table 8).

Incase of interaction effect, data was recorded between nitrogen and phosphorous in consideration of fruit length under the present experiment and found statistically significant variation (Appendix V). The maximum fruit length (5.610 cm) was recorded from $N_{150}P_{150}$ treatment consisting of 150 kg N/ha +100 kg P_2O_5 /ha, and the minimum fruit length per plant (4.52 cm) was recorded from N_0P_0 treatment (Table 9) and founds with N_0P_{150} , N_0P_{100} and N_0P_{150} treatments. Islam *et al.* (1997) showed that the length and breadth of individual fruit increased with increasing of nitrogen level.

4.1.11 Fruit size (Fruit breadth)

Breadth of individual fruit per plant incase of fruit size varied significantly due to the application of nitrogen in tomato (Appendix V). The highest fruit breadth (5.88 cm) was found from N_{200} treatment consisting of 200 kg N/ha, which was statistically identical (5.80 cm) with both N_{150} and N_{100} treatment consisting of 150 kg N/ha, and 100 kg N/ha. And the lowest fruit breadth (4.283 cm) was recorded from N_0 treatment (Table 8). The ranges of Fruit breadth incase of nitrogen application were 4.283 cm to 5.88 cm.

In regard of different level of phosphorous data showed statistically significant difference for fruit breadth per plant (Appendix V). The maximum fruit breadth (4.6 cm) was observed from P_{150} treatment consisting of 150 kg P_2O_5 /ha which

was statistically identical (4.568 cm) with P_{100} treatment as 100 kg P_2O_5 /ha while (4.505 cm) with P_{50} treatment as 50 kg P_2O_5 /ha and the minimum fruit breadth (4.358 cm) was recorded from P_0 treatment (Table 8) The ranges of Fruit breadth incase of phosphorous application were 4.358 cm to 4.600 cm.

Due to interaction effect was also recorded between nitrogen and phosphorous in consideration of fruit breadth per plant and also found statistically significant variation (Appendix V). The maximum fruit breadth per plant (4.700cm) was recorded from both $N_{100}P_{100}$ and $N_{150}P_{100}$ treatment consisting of 100 kg N/ha +100 kg P_2O_5 /ha and $N_{150}P_{100}$ treatment consisting of 150 kg N/ha +100 kg P_2O_5 /ha and on the other hand the minimum fruit breadth (4.11cm) was recorded from N_0P_0 treatment (Table 9) where no fertilizers were applied.

4.1.12 Fruit weight per plant

Weight of fruit per plant varied significantly due to the application of different doses of nitrogen in tomato (Appendix V).The maximum weight fruit fruits per plant (2.125 kg) was recorded from N_{200} treatment consisting of 200 kg N/ha, while the minimum weight of fruit per plant (1.085kg) was observed from N_0 treatment (Table 8). The ranges of fruit weight per plant incase of nitrogen application were 1.085 kg to 2.125 kg. Chung *et al.* (1992) reported that increasing levels of nitrogen increased the fresh weight of tomato fruit.

Due to different level of phosphorous application data showed statistically significant difference for weight of fruit per plant (Appendix V). The maximum weight of fruit per plant (2.033 kg) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical (2.007 kg) with P_{100} treatment as 100 kg P_2O_5 /ha and where as the minimum weight of fruits per plant (1.225 kg) was recorded from P_0 treatment (Table 8). Ahmed and Choudhury (1990) reported that with increasing the level of phosphorous, plant growth increases and which ensure maximum number of flower and fruits per cluster as well as the weight of fruits.

Table 8. Effect of nitrogen and phosphorous on fruit size, weight of individual fruit, fruit weight /plant and fruit yield of tomato

Treatment	Fruit size		Weight of individual fruit (gm)	Fruit weight /plant (kg)	Fruit yield (ton/ha)
	Length (cm)	Breadth (cm)			
Nitrogen					
N ₀	4.605b	4.289b	66.22c	1.085d	39.00d
N ₁₀₀	5.225a	4.580a	80.40b	1.690c	60.91c
N ₁₅₀	5.320a	4.580a	83.77a	2.050b	73.80b
N ₂₀₀	5.278a	4.588a	83.27a	2.125a	76.52a
LSD	0.114	0.087	2.121	0.058	1.919
Phosphorous					
P ₀	4.960c	4.358b	71.54b	1.225c	44.11c
P ₅₀	5.080bc	4.505a	79.49a	1.685b	60.62b
P ₁₀₀	5.125b	4.568a	80.96a	2.007a	72.24a
P ₁₅₀	5.262a	4.600a	81.67a	2.033a	73.26a
LSD	0.132	0.101	2.449	0.068	2.216

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability.

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Table 9. Interaction effect of nitrogen and phosphorous on fruit size, weight of individual fruit, fruit weight /plant and fruit yield of tomato

Treatments (N×P)	Fruit size		Weight of individual fruit (gm)	Fruit weight /plant (kg)	Fruit yield (ton/ha)
	Length (cm)	Breadth (cm)			
N ₀ P ₀	4.520e	4.110f	63.85d	0.7700i	27.72j
N ₀ P ₅₀	4.600e	4.300e	67.37d	1.080h	38.83i
N ₀ P ₁₀₀	4.650e	4.320de	65.99d	1.220g	43.78h
N ₀ P ₁₅₀	4.650e	4.400de	67.68d	1.270g	45.67h
N ₁₀₀ P ₀	4.900d	4.420cde	73.03c	1.220g	43.79h
N ₁₀₀ P ₅₀	5.200c	4.500abcd	80.86b	1.680d	60.50e
N ₁₀₀ P ₁₀₀	5.300bc	4.700a	83.31ab	1.870c	67.48d
N ₁₀₀ P ₁₅₀	5.500ab	4.700a	84.41ab	1.990bc	71.88bc
N ₁₅₀ P ₀	5.200c	4.450bcde	74.87c	1.390f	50.07g
N ₁₅₀ P ₅₀	5.220c	4.620ab	85.32ab	1.930bc	69.48cd
N ₁₅₀ P ₁₀₀	5.250bc	4.600abc	87.59a	2.440a	87.76a
N ₁₅₀ P ₁₅₀	5.610a	4.650ab	87.29a	2.440a	87.89a
N ₂₀₀ P ₀	5.220c	4.450bcde	74.41c	1.520e	54.87f
N ₂₀₀ P ₅₀	5.300bc	4.600abc	84.42ab	2.050b	73.67b
N ₂₀₀ P ₁₀₀	5.300bc	4.650ab	86.97a	2.500a	89.93a
N ₂₀₀ P ₁₅₀	5.290bc	4.650ab	87.29a	2.430a	87.60a
LSD	0.2299	0.1749	4.242	0.1179	3.839
CV (%)	2.67	2.36	3.24	3.98	3.68

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability.

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Interaction effect was also recorded between nitrogen and phosphorous in consideration of weight of fruits per plant and found statistically significant variation (Appendix V). The highest weight of fruits per plant (2.5 kg) was recorded from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha +100 kg P_2O_5 /ha which was statistically identical (2.44 kg), with the treatments $N_{150}P_{100}$ and $N_{150}P_{150}$ consisting of 150 kg N/ha +100 kg P_2O_5 /ha and 150 kg N/ha +150 kg P_2O_5 /ha and (2.430 kg) from $N_{200}P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha. While the lowest weight of fruits per plant (0.770 kg) was recorded from N_0P_0 treatment (Table 9). Ahmed and Choudhury (1990) revealed that higher dose of nitrogen and phosphorous is influential nutrient for attaining better growth and ultimate results was desirable growth with maximum number of flower and fruit per cluster and highest weight of fruits per plant.

4.1.13 Fruit yield

Fruit yield of tomato varied significantly due to application of nitrogen in tomato (Appendix V). The maximum fruit yield (76.72 ton/ha) (Figure 2) was recorded from N_{200} treatment consisting of 200 kg N/ha, and the minimum fruit yield (39.0 ton/ha) was recorded from N_0 treatment (Table 8). Chung *et al.* (1992) reported that increasing levels of nitrogen increased the fresh weight of tomato fruit as well as fruit yield.

Different level of phosphorous showed statistically significant difference for fruit yield in tomato (Appendix V). The maximum fruit yield (73.26 ton/ha) was found from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical (72.24 ton/ha) (Figure 3) with P_{100} treatment as 100 kg P_2O_5 /ha and while the minimum fruit yield (44.11 ton/ha) was recorded from P_0 treatment (Table 8). With increasing the level of phosphorous, plant growth increases and which ensure maximum number of flower and fruits per cluster as well as the weight of fruits and also increase the fruit yield of tomato.




Figure 2. Effect of nitrogen on the yield (ton/ha) of tomato




Figure 3. Effect of phosphorous on the yield (ton/ha) of tomato




Figure 4. Interaction effect of nitrogen and phosphorous on the yield (ton/ha) of tomato

Interaction effect was also recorded between nitrogen and phosphorous in consideration of fruit yield of tomato and found statistically significant variation (Appendix V). The maximum fruit yield (89.93 ton/ha) was recorded from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha +100 kg P_2O_5 /ha (Figure 4) which was statistically identical $N_{150}P_{150}$ (87.89 ton/ha) $N_{150}P_{100}$ (87.76 ton/ha) and $N_{200}P_{150}$ (87.60 ton/ha) treatment consisting of 150 kg N/ha +150 kg P_2O_5 /ha, 150 kg N/ha +100 kg P_2O_5 /ha and 200 kg N/ha +150 kg P_2O_5 /ha, respectively. The minimum fruit yield (27.72 ton/ha) was recorded from N_0P_0 treatment (Table 9).

4.2 Nutrient content in shoot

The shoot nitrogen and phosphorous concentration were determined for different level of nitrogen and phosphorous application and also their combination.

4.2.1 Nitrogen content in shoot

Application of different level of nitrogen showed a statistically significant variation for nitrogen content in shoot of tomato (Appendix VII). application of N increased the N content in shoot and the highest concentration shoot N found with in highest dose of N. The maximum nitrogen content (1.742%) was recorded from N_{200} treatment consisting of 200 kg N/ha, which was closely followed by N_{150} (1.487%) treatment as 150 kg N/ha and the minimum nitrogen content (0.74%) was recorded from N_0 treatment, which was closely (1.357%) followed by N_{100} treatment as 100 kg N/ha (Table 10). The ranges of Nitrogen content (%) in shoot incase of nitrogen application were 0.74% to 1.742%. Soares *et al.*, (1986) reported that N fertilization increased N content in the shoot and fruits. Velenzuela *et al.*, (1987) also reported that straw N content was increased with increasing rates of applied N.

Due to application of different level of phosphorous data showed statistically significant differences for Nitrogen content in tomato shoot or straw (Appendix VII). The maximum nitrogen content (1.497%), was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical (1.423%) with P_{100} treatment as 100 kg P_2O_5 /ha and the minimum nitrogen content (1.115%) was

recorded from P_0 treatment (Table 10). The ranges of nitrogen content (%) in shoot incase of phosphorous application were 1.115% to 1.497%.

In regard to interaction effect was also recorded between nitrogen and phosphorous in consideration of nitrogen content in tomato shoot or straw and found statistically significant variation (Appendix VII). The maximum nitrogen content in tomato shoot or straw (2.040%) was recorded from $N_{200}P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha. The minimum nitrogen content in tomato shoot or straw (0.50%) was recorded from N_0P_0 treatment (Table 11).

4.2.2 Phosphorous content (ppm) in shoot

Effect of different level of nitrogen showed a statistically significant variation for phosphorous content (ppm) in straw or shoot of tomato (Appendix VII). The ranges of Phosphorous content (ppm) in straw incase of nitrogen application were 32.25 ppm to 55.98 ppm. The maximum phosphorous content (55.98 ppm) was recorded from N_{200} treatment consisting of 200 kg N/ha, which was closely (50.88 ppm) followed by N_{150} treatment as 150 kg N/ha and the minimum phosphorous content (32.25 ppm) was recorded from N_0 treatment (Table 10).

Application of different level of phosphorous showed statistically significant difference for phosphorous content in tomato shoot or straw (Appendix VII). The maximum phosphorous content (55.80 ppm) was observed from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was closely (52.48 ppm) followed by P_{100} treatment as 100 kg P_2O_5 /ha, and the minimum phosphorous content (28.80 ppm) was recorded from P_0 treatment (Table 10). Wateren and Coltman (1988) reported that increasing P levels increased shoot P concentrations. Dhinakaran and Savitri (1997) found that P content of tomato fruit increased with increased application of P.

Table 10. Effect of nitrogen and phosphorous application on nitrogen & phosphorous in shoot and fruit of tomato

Treatments	N (%) in shoot	P (ppm) in shoot	N % in fruit	P % in fruit
Nitrogen				
N ₀	0.740d	32.25d	2.852c	0.200c
N ₁₀₀	1.357c	40.65c	3.257b	0.227b
N ₁₅₀	1.457b	50.88b	3.600a	0.247a
N ₂₀₀	1.742a	55.98a	3.662a	0.240ab
LSD	0.074	1.947	0.131	0.0186
Phosphorous				
P ₀	1.115c	28.80d	3.155c	0.190c
P ₅₀	1.293b	42.67c	3.280bc	0.225b
P ₁₀₀	1.423a	52.48b	3.425ab	0.245ab
P ₁₅₀	1.497a	55.80a	3.513a	0.255a
LSD	0.086	2.249	0.152	0.0215

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Table 11. Interaction effect of nitrogen and phosphorous application on nitrogen & phosphorous in shoot and fruit of tomato

Treatments (N×P)	Nitrogen (%) in shoot	Phosphorous (ppm) in shoot	N % in fruit	P % in fruit
N ₀ P ₀	0.50i	25.00h	2.650i	0.170f
N ₀ P ₅₀	0.70h	30.00g	2.860hi	0.200def
N ₀ P ₁₀₀	0.88g	36.00f	2.900ghi	0.210def
N ₀ P ₁₅₀	0.88g	38.00f	3.000fgh	0.220cde
N ₁₀₀ P ₀	1.26f	27.10gh	3.170efg	0.200def
N ₁₀₀ P ₅₀	1.31f	35.30f	3.200def	0.230bcde
N ₁₀₀ P ₁₀₀	1.37ef	50.00e	3.260def	0.240abcd
N ₁₀₀ P ₁₅₀	1.49ce	50.20e	3.400cde	0.240abcd
N ₁₅₀ P ₀	1.30f	28.00gh	3.370cde	0.200def
N ₁₅₀ P ₅₀	1.52ce	50.20e	3.490bcd	0.240abcd
N ₁₅₀ P ₁₀₀	1.55cd	60.30c	3.740ab	0.270ab
N ₁₅₀ P ₁₅₀	1.58c	65.00b	3.800a	0.280a
N ₂₀₀ P ₀	1.40def	35.10f	3.430cde	0.190ef
N ₂₀₀ P ₅₀	1.64c	55.20d	3.570abc	0.230bcde
N ₂₀₀ P ₁₀₀	1.89b	63.63bc	3.800a	0.260abc
N ₂₀₀ P ₁₅₀	2.040a	70.00a	3.850a	0.280a
LSD	0.1491	3.895	0.2637	0.0372
CV (%)	6.86	5.20	4.72	6.72

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha

Due to interaction effect, data was recorded between nitrogen and phosphorous in consideration of phosphorous content in tomato shoot or straw and found statistically significant variation (Appendix VII). The highest amount of phosphorous content in tomato shoot or straw (70.00 ppm) was recorded from $N_{200}P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha. The lowest amount of phosphorous content in tomato shoot or straw (25.00 ppm) was recorded from N_0P_0 treatment that is control condition (Table 11).

4.3 Nutrient content in fruit

Nutrient such as nitrogen, phosphorous and potassium content in fruit was estimated for different level of nitrogen and phosphorous application and also their combination.

4.3.1 Nitrogen content (%) in fruit

Due to application of different level of nitrogen showed a statistically significant variation for nitrogen content in fruit of tomato (Appendix VII). The ranges of nitrogen content (%) in fruit incase of nitrogen application were 2.852 % to 3.662 %. The maximum nitrogen content (3.662%) was recorded from N_{200} treatment consisting of 200 kg N/ha, which was statistically identical (3.600%) with N_{150} treatment as 150 kg N /ha, and the minimum nitrogen content (2.852%) was recorded from N_0 treatment (Table 10).

Different level of phosphorous showed statistically significant difference for nitrogen content in tomato fruit (Appendix VII). The ranges of nitrogen content (%) in fruit incase of phosphorous application were 3.155 % to 3.513 %. The maximum nitrogen content (3.513) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha, and the minimum nitrogen content (3.155%) was recorded from P_0 treatment (Table 10).



In respect of interaction effect was also recorded between nitrogen and phosphorous in consideration of nitrogen content in tomato fruit and found statistically significant variation (Appendix VII). The maximum nitrogen content in tomato fruit (3.850%) was recorded from $N_{200}P_{150}$ treatment consisting of 200 kg N/ha + 150 kg P_2O_5 /ha, which was statistically identical in both $N_{200}P_{100}$ (3.8%) and $N_{150}P_{150}$ (3.8%) treatment consisting of 200 kg N/ha + 100 kg P_2O_5 /ha and 150 kg N/ha + 150 kg P_2O_5 /ha respectively. The minimum nitrogen content in tomato fruit (2.650%) was recorded from N_0P_0 treatment that is control condition (Table 11).

4.3.2 Phosphorous content (%) in fruit:

Effect of different level of nitrogen showed a statistically significant variation for phosphorous content (%) in fruit of tomato under the present trial (Appendix VII). The ranges of phosphorous content (%) in fruit incase of phosphorous application were 0.200% to 0.2475%. The maximum phosphorous content (0.2475%) was recorded from N_{150} treatment consisting of 150 kg N/ha, which was closely (0.2400%) followed by N_{200} treatment as 200 kg N/ha and the minimum phosphorous content (0.2%) was recorded from N_0 treatment (Table 10).

Application of different level of phosphorous showed statistically significant difference for phosphorous content in tomato fruit (Appendix VII). On the one hand, the maximum phosphorous content (0.2550%), was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was closely (0.2450%) followed by P_{100} treatment as 100 kg P_2O_5 /ha, and on the other hand, the minimum phosphorous content (0.1900%) was recorded from P_0 treatment that is control condition (Table 10).

Due to interaction effect was also recorded between nitrogen and phosphorous in consideration of phosphorous content in tomato fruit and found statistically significant variation (Appendix VII). The highest amount of phosphorous content in tomato fruit (0.280%) was recorded from both $N_{200}P_{150}$ and $N_{150}P_{150}$ treatment

consisting of 200 kg N/ha +150 kg P_2O_5 /ha and 150 kg N/ha +150 kg P_2O_5 /ha. The lowest amount of phosphorous content in tomato fruit (0.1700%) was recorded from N_0P_0 treatment that is control condition (Table 11).

4.4 Status of post harvest soil

4.4.1 pH at post harvest soil

Effect of different level of nitrogen showed a statistically non significant for pH at post harvest soil under the present trial (Appendix VI). The highest pH content (5.4) was recorded from N_{100} treatment consisting of 100 kg N/ha, and the lowest pH content (5.3) was recorded from N_0 treatment (Table 12).

Application of different level of phosphorous also showed statistically non significant for pH at post harvest soil (Appendix VI). The maximum pH content (5.425) was recorded from P_{100} treatment consisting of 100 kg P_2O_5 /ha, and the minimum pH content (5.25) was recorded from P_0 treatment (Table 12)

Interaction effect was also recorded between nitrogen and phosphorous in consideration of P^H at post harvest soil under the present experiment and found statistically non significant (Appendix VI). The maximum pH content (5.6) was recorded from both $N_{100}P_{100}$ and $N_{200}P_{100}$ treatments consisting of 100 kg N/ha +100 kg P_2O_5 /ha and 200 kg N/ha +100 kg P_2O_5 /ha and 100 kg N/ha +150 kg P_2O_5 /ha. The minimum pH content (5.0) was recorded from N_0P_0 treatment (Table 13). Due to application of nitrogen and phosphorous in different doses for tomato the pH of the post harvest soil has increased slightly over the initial soil pH. The initial soil pH of the soil was 5.0 and the pH became 5.00 to 5.60 that is slightly increasing pattern.

4.4.2 Nitrogen at post harvest soil

Application of different level of nitrogen showed a statistically significant variation for nitrogen content (%) at post harvest soil (Appendix VI). The maximum nitrogen content (0.114%) was recorded from N_{200} treatment consisting of 200 kg N/ha, which was statistically identical with N_{150} (0.1123%) treatment as 150 kg N/ha and was recorded from N_{100} (0.1107%) treatment as 100 kg N/ha, and

the minimum nitrogen content (0.097%) was recorded from N_0 treatment (Table 12). The ranges of nitrogen (%) at post harvest soil in case of nitrogen application were 0.097% to 0.114%. Ahmed (1992) reported that highest application of nitrogen accelerated nitrogen uptake by soil and the ultimately result is the highest nitrogen content in soil.

Due to application of different level of phosphorous showed statistically non significant for nitrogen content (%) at post harvest soil (Appendix VI). The maximum nitrogen content (0.1125%) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha which was statistically identical (0.1105%) with P_{100} treatment as 100 kg P_2O_5 /ha and (0.1070%) P_{50} treatment as 50 kg P_2O_5 /ha, and the minimum nitrogen content (0.1042%) was recorded from P_0 treatment that is control condition (Table 12). The ranges of nitrogen (%) at post harvest soil in case of phosphorous application were 0.1042% to 0.1125%.

In regard to interaction effect was also recorded between nitrogen and phosphorous in consideration of nitrogen (%) at post harvest soil and found statistically non significant similar (Appendix VI). The maximum nitrogen (%) at post harvest soil (0.119%) was recorded from $N_{200}P_{150}$ treatment consisting of 200 kg N/ha + 150 kg P_2O_5 /ha. The minimum nitrogen content (0.096%) was recorded from N_0P_0 treatment (Table 13).

Table 12. Effect of nitrogen and phosphorous on pH, nitrogen and phosphorous of post harvest soil

Treatments	pH	N (%)	P (ppm)
Nitrogen			
N ₀	5.25	0.09b	27.68c
N ₁₀₀	5.42	0.11a	32.24b
N ₁₅₀	5.30	0.11a	36.34a
N ₂₀₀	5.37	0.11a	36.85a
LSD	NS	0.01318	1.692
Phosphorous			
P ₀	5.25	0.104	22.25d
P ₅₀	5.27	0.107	30.89c
P ₁₀₀	5.42	0.110	37.82b
P ₁₅₀	5.4	0.112	42.15a
LSD	NS	NS	1.953

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha

P₁₅₀: 150 kg P₂O₅ /ha



Table 13. Interaction effect of nitrogen and phosphorous on pH, nitrogen concentration and phosphorous concentration of post harvest soil

Treatments	pH	N (%)	P (ppm)
N ₀ P ₀	5.0	0.098	13.84g
N ₀ P ₅₀	5.3	0.096	26.32ef
N ₀ P ₁₀₀	5.2	0.097	34.54c
N ₀ P ₁₅₀	5.5	0.097	36.01c
N ₁₀₀ P ₀	5.5	0.104	23.31f
N ₁₀₀ P ₅₀	5.3	0.110	29.07de
N ₁₀₀ P ₁₀₀	5.6	0.113	34.07c
N ₁₀₀ P ₁₅₀	5.3	0.116	42.52ab
N ₁₅₀ P ₀	5.4	0.107	26.00ef
N ₁₅₀ P ₅₀	5.1	0.108	32.39cd
N ₁₅₀ P ₁₀₀	5.3	0.116	41.67ab
N ₁₅₀ P ₁₅₀	5.4	0.118	41.01b
N ₂₀₀ P ₀	5.1	0.108	25.84ef
N ₂₀₀ P ₅₀	5.4	0.114	35.77c
N ₂₀₀ P ₁₀₀	5.6	0.116	45.30a
N ₂₀₀ P ₁₅₀	5.4	0.119	44.77a
LSD	NS	NS	3.383
CV	3.65	3.89	6.10

In a column means having common letter(s) are statistically identical and those having dissimilar letter(s) differ significantly as per 0.01 level of probability

N₀: 0 kg N/ha

N₁₀₀: 100 kg N/ha

N₁₅₀: 150 kg N/ha

N₂₀₀: 200 kg N/ha

P₀: 0 kg P₂O₅ /ha

P₅₀: 50 kg P₂O₅ /ha

P₁₀₀: 100 kg P₂O₅ /ha


P₁₅₀: 150 kg P₂O₅ /ha

4.4.3 Phosphorous content at post harvest soil

Effect of different level of nitrogen showed a statistically significant variation for phosphorous content (ppm) at post harvest soil (Appendix VI). The maximum phosphorous content (36.85 ppm) was recorded from P_{200} treatment consisting of 200 kg N/ha, (Figure 23) which was statistically identical and (36.34 ppm) was recorded from N_{150} treatment as 150 kg N/ha and the minimum phosphorous content (17.68 ppm) was recorded from N_0 treatment (Table 12). The ranges of phosphorous content (ppm) at post harvest soil incase of nitrogen application were 17.68 ppm to 36.85 ppm.

Application of different level of phosphorous showed statistically significant variation for phosphorous content (ppm) at post harvest soil (Appendix VI). The maximum phosphorous content (42.15 ppm) was recorded from P_{150} treatment consisting of 150 kg P_2O_5 /ha and the minimum phosphorous content (22.25 ppm) was recorded from P_0 treatment (Table 12). The ranges of phosphorous content (ppm) at post harvest soil incase of phosphorous application were 22.25 ppm to 42.15 ppm.

Due to interaction effect was also recorded between nitrogen and phosphorous in consideration of phosphorous (ppm) at post harvest soil and found statistically significant variation (Appendix VI). The maximum phosphorous (45.30 ppm) at post harvest soil was recorded from $N_{200}P_{100}$ treatment consisting of 200 kg N/ha +100 kg P_2O_5 /ha which was statistically identical (44.77 ppm) at post harvest soil was recorded from $N_{200}P_{150}$ treatment consisting of 200 kg N/ha +150 kg P_2O_5 /ha. The minimum phosphorous content (13.84 ppm) was recorded from N_0P_0 treatment (Table 13).



Chapter V

Summary and Conclusion

CHAPTER V

SUMMARY AND CONCLUSION

The experiment was conducted at the Research Farm of Sher-e-Bangla Agricultural University, Sher-e Bangla Nagar, Dhaka, during November 2008 to March 2009 to find out the effect of nitrogen (N) and phosphorous (P) on the growth and yield of BARI Tomato-2 which is one of the high yielding varieties of indeterminate type. BARI Tomato-2 (Raton), a high yielding variety of tomato (*Lycopersicon esculentum* Mill.) developed by Bangladesh Agricultural Research Institute (BARI), Gazipur. The seeds were collected from Horticultural Research Centre (HRC) of BARI, Gazipur. Healthy and uniform sized 30 days old seedlings were uprooted from the seed bed and were transplanted in the experimental field on 28 November, 2008 maintaining a spacing of 60 cm and 50 cm between the rows and plants, respectively. The experiment was laid out in a Randomized Complete Block Design (RCBD) with three replications of each fertilizer treatment combinations. Fertilizer treatments consisting of 4 levels of N and 4 levels of P on the growth and yield of tomato. The study comprised the following treatments: Four levels of nitrogen N_0 : 0 kg N/ha, N_{100} : 100 kg N/ha, N_{150} : 150 kg N/ha and N_{200} : 200 kg N/ha and Four levels of phosphorous: P_0 : 0 kg P_2O_5 /ha, P_{50} : 50 kg P_2O_5 /ha, P_{100} : 100 kg P_2O_5 /ha and P_{150} : 150 kg P_2O_5 /ha. There were 16 (4 × 4) treatment combinations. Data were recorded on yield and yield contributing characters and yield of tomato and nitrogen and phosphorus uptake by plant and their availability on soil.


Yield and yield contributing characters, N and P concentration in plant and post harvest soil were recorded. Plant height at 60 days after transplanting, number of branches per plant, number of flowers per plant, number of fruits of plant and fruit weight per plant increased significantly with increasing N level up to 150 kg N/ha, whereas fruit yield increased significantly up to 200 kg N/ha. All these parameters also increased significantly with increasing the level of P up to 150 kg P/ha.



However, the treatment combination $N_{200}P_{150}$ showed the maximum plant height at harvest (93.62 cm). On the other hand, the treatment combination $N_{200}P_{100}$ produced the maximum number of flowers per plant (49.86), number of fruits per cluster (3.99), number of fruits per plant (38.30) fruit weight (2.5 kg/plant) and fruit yield (89.93 t/ha). N and P concentrations in shoot and fruit were the highest in $N_{200}P_{150}$ treatment. The treatment had no significant effect on pH of post harvest soil. However increasing the application of nitrogen increased soil total N and P content up to N_{150} treatment. Application of P had not significant effect on total nitrogen but P_{150} produced the highest P content in soil. $N_{200}P_{150}$ combination had the highest N (0.119%) and P (44.77 ppm) concentration in plant.

Considering the situation of the present experiment, further studies in the following areas may be suggested:

1. The treatment combination $N_{200} P_{100}$ can be recommended for Tomato cultivation in this region.
2. Such Study is needed in various AEZs with different Tomato varieties to find out the regional adaptability performances.
3. Another level of N may be included in this experimental protocol.



Chapter References

REFERENCES

- Adjanohoun, A., Hernandez, J. A. and Berenguer, T. 1996. Response of tomato (*Lycopersicon esculentum*, Mill.) to nitrogen fertigation on a ferrallitic red soil. *Cultivos Tropicales*. 17(2): 23-24.
- Ahmed, S. 1992. Crops response to applied nitrogen and phosphorus in Brahmaputra flood plain soil of Bangladesh. *Bangladesh J. Soil Sci.*, 19:36-41.
- Ahmed, C.M.S. and M.A. Choudhury, 1990. Effect of nitrogen on some yields parameters in tomato (*Lycopersicon esculentum*, Mill.). Under Haidugari environmental condations. *Sarad J. of Agriculture* 6(2): 155-158.
- Al-Afifi, M.A. Itani, S. haris, H. Al-Masri, IA. Al-Gharib and S. Khalil. 1993. Response of processing tomato to nitrogen and phosphorus application at moderate low temperature. *Egyptian J. Hort.*, 18(1): 45-62.
- Annanurova, M.A.; M. Rozyeva, T. Tailakov and L.P. Slovinskaya. 1992. Effect of fertilizers on some physiological process and fruit quality in tomatoes. *Izvestiya Akademii Nauk Turkmenistan* 3: 49-52.
- Anonymous, 2008. Annual Weather Report, Meteorological Station, Dhaka. Bangladesh.
- Awad, S. S., Hassan, H. M. F., Shahien, A. H. and Zayed, A. A. 2001. Studies on intercropping of parsley and demsisa with tomato under different rates of nitrogen fertilization. *Alexandria J. Agril. Res.*, 46(2): 97- 112.
- Banerjee, M. K., Balyan, D. S., Kalloo, G., Sing, A. and Saini, P. S. 1997. Effect of nitrogen fertilization and planting pattern on fruit yield of tomato cv. Hisar Lalima. *Crop Res.*, 14(3): 441-446.
- BARC. 2005. Fertilizer recommendation guide. Bangladesh Agricultural Research Council. Farmgate, Dhaka-1215, pp. 1-72.
- Bar-Yosef, B., Imas, P. Adams, P. Hidding, AP. Kipp-JA.C. Sonneveld, and C. kreij. 1995. Production and nutrient contents in greenhouse tomatos. *Acta Hort.*, 401 :337-346.

- BBS. 2006, Yearly Statistical Book. Bangladesh Bureau of Statistics, Dhaka, Bangladesh. 43p.
- Belichki, I. 1984. Fertilization of the tomato cultivar Triump grown for early field production. *Gard insrska*, 20(2): 57-63.
- Bellert, C., Bot, J., Dorais, M., Lopez, J., Gosselin, A. Bot, J. and Munoz, C. R. 1998. Nitrogen accumulation and growth of fruiting tomato plants in hydroponics. *Acta Horticulturae*. 458: 293-301.
- Lose, T. K. and Som, M. G. 1990. Vegetable crops in India. Naya Prakash, Calcutta, India. pp. 687-691.
- Bot, J., Jeannequin, B., Fabre, R. and Bot, J. 2001. Growth and nitrogen status of soilless tomato plants following nitrate withdrawal from the nutrient solution. *Annals Bot.*, **88**: 3, 361-370.
- Bremner, J. M. and Mulvaney, C. S. 1982. Total Nitrogen. In Methods of Soil Analysis. Miller, R. H. and Xeemy, D. R. 1982. Amer. Soc. Agron. Inc. Madi. Wis. USA. pp.593-622.
- Candilo, M. D. L., Leoni, C. and Silvestri, G. P. 1993. Sulphur, phosphorus and potassium in processing tomato in high alkaline soil. *Adv. Hort.*, 7(2): 57-60.
- Cerne, M. and L. Briski. 1993. Nutrition and irrigation of tomato. *Acta Hort.*, **376**:319, 322.
- Ceylan, S., Mordogan, N., Yoldas, F. and Yagmur, B. 2001. The effect of nitrogen fertilization on yield, nitrogen accumulation and content of nutrients in the tomato plant. *Ege Universitie Ziraat Fakultesi Dergisi*. **38**(2-3): 103-110.
- Chang, K., Sommer, K. and Chang, K. R. 2001. The effect of nitrogen supply to tomato plants by NH_4 beaker deposits. *J. the Korean Soc. Hort. Sci.*, **41**(4): 334-338.
- Christian, G. D. and F. J. Feldman. 1970. Atomic absorption spectroscopy, Application in Agriculture, Biology and Medicine, Wiley-Inter-science. New York.



- Choudhury, B. 1997. Vegetables (6th Revised Edn). The Director, National Book Trust. New Delhi, India. P. 46.
- Chung, S. J., Seo, B. S. and Lee, B. S. 1992. Effects of nitrogen, potassium levels and their interaction on the growth and development of hydroponically grown tomato. *J Kor. Soc. Hort. Sci.*, **33**(3): 244-251.
- Cuortero, J. and Fernandez, R. 1999. Tomato and salinity. *Scientia Hort.*, **78**(1-4): 83-84.
- Dufault, R. J., Decoteau, D. R., Garrett, J. T., Batal, K. D., Granberry, D., Davis, J. M., Hoyt, G. and Sanders, D. 2000. Influence of cover crops and inorganic nitrogen fertilization on tomato and snap bean production and soil nitrate distribution. *J. Veg. Crop Prod.*, **6**(2): 13-25.
- FAO. 1988. Production Year Book. Food and Agriculture Organization. Rome, Italy. 42: 190-193.
- FAO. 1997. FAO Production Year Book. Basic Data Unit. Statistics Division, FAO. Rome, Italy, 51: 125-127.
- Felipe, E .F. and Casanova, O. E. 2000. Nitrogen, phosphorus and potassium fertilization in tomato (*Lycopersicon esculentum* Mill.) in alluvial bank soils of the Guarico river. *Reivista Unellez de Cienciay Tecnologia, Production Agricola*. **17**(1): 21-44.
- Garton, R. W. and Widders, I. E. 1990. Nitrogen and phosphorus preconditioning of small- plug seedlings influence processing tomato productivity. *Hort. Science*. **25**(6):655-657.
- Gomez, K. A. and Gomez, A. A. 1984. Statistical Procedure for Agricultural Research (2nd edn.). Int Rice Res. Inst., *A Willey Int. Sci.*, Pub., pp. 28-192.
- Grela, L. M., Delgado, N. M., Jimenez, R. R Huerres, P. C. and Grela, L. H. 1988. Effect of different nitrogen rates and plant spacing on growth and development of commercial tomato (*Lycopersicon esculentum* Mill.) cultivars. *Cetro-Agricola*, **15**(4): 55-62.

- Groot, C. C., Marcelis, L. F. M. 2001. Regulation of growth by phosphorus supply in whole tomato plants. Plant nutrition food security and sustainability of agro ecosystems through basic and applied research. Fourteenth International Plant Nutrition Colloquium, Hannover, Germany. 114-115.
- Gupta, A. and Shukla, V. 1977. Response of tomato (*Lycopersicon esculentum* Mill) to plant spacing, nitrogen, phosphorous and potassium fertilization. *Indian J. Hort.*, **33**:270-276.
- Gupta, C. R, Sengar, S. S. 2001. Response of tomato (*Lycopersicon esculentum* Mill.) to nitrogen and potassium fertilization in acidic soil of Bastar. *Veget. Sci.* **27**(1): 94-95.
- Hedge, D. M. 1997. Nutrient requirements of solanaceous vegetable crops. Food and Fertilizer Technology Centre for the Asia and Pacific Region Extension Bulletin, Taipei, 441:9.
- Hoffland, E. Dicke, M., Tintelen, W. Dijkman, H. and Beusichem, M. L. 2000. Nitrogen availability and defense of tomato against two-spotted spider mite. *J. Chemical Ecol.*, **26**(12):2697-2711.
- Hoffland, E., Beusichem, M. L., Jeger, M. J. and Beusichem, M. L. 1999. Nitrogen availability and susceptibility of tomato leaves to *Botrytis cinerea*. *Plant and Soil.* **210**(2): 263-272.
- Hohjo, M., Kuwata, C., and Yoshikawa, K., and Tognoni, F., Namiki, T., Nukaya, A. and Maruo, T. 1995. Effects of nitrogen form, nutrient concentration and Ca concentration on the growth, yield and fruit quality in NFT-tomato plants. *Acta Hort.*, **396**: 145-152.
- Hunter, A.H. 1980. "Soil Fertility Analytical Service in Bangladesh" consultancy report, BARI Project Phase II, BARC, Dhaka, Bangladesh.
- Islam, M.A., A.M. Farooque, A. Siddiqua and A. Siddque. 1997. Effect of planting patterns and different nitrogen levels on yield and quality of tomato. *Bangladesh J. Agril. Sci.*, **24**(1): 9-15

- Kaya, C. and Higgs, D. 2001. Growth enhancement by supplementary phosphorus and iron in tomato cultivars grown hydroponically at high zinc. *J. Plant Nutri.*, **24**(12): 1861-1870.
- Kaya, C., Kirnak, H. and Higgs, D. 2001. Enhancement of growth and normal growth parameters by foliar application of potassium and phosphorus in tomato cultivars grown at high salinity. *J. Plant Nutri.*, **24**(2):357-367.
- Khalil, S. A. Noor. B., Kausar. M. A., Muhammad. A. Shah, S. A. 2001. Response of tomato to different nitrogen fertilizers alone and in combination with phosphorus and potassium. *Sarhad J. Agril.*, **17**(2): 213-217.
- Kooner, K. S. and Randhawa, K. S. 1990. Effect of varying levels and sources of nitrogen on yield and processing qualities of tomato varieties. *Acta-Horticulturae*. 267: 93-99; Proceedings of the 6th symposium on the timing of field production of vegetables, held at Wageningen, Netherlands, 21-25 Aug. 1989.
- Kuksal, R. P., R.D. Sign and M.R. Yadav. 1977. Effect of different levels of nitrogen and phosphorus on fruit and seed of tomato variety Chaubatti Red. *Prog. Hort.*, **9**(2): 13- 20.
- Manang, E. Z., Uriyo, A. P. and Singh, B. R. 1982. Effects of fertilizer nitrogen and phosphorous on tomato. *Beitriige zur tropischen land wirtschaft and veterinar medizin Dares Salaam University, Morogoro, Tanzania*, **20**(3): 247-253.
- Manoj, R. and Raghav, M. 2001. Effect of nitrogen and spacing on growth and yield of hybrid tomato. *Advances in Horticulture and Forestry*. **8**: 103-107.
- Melton, R. C. and Dufault, R. S. 1991. Nitrogen, phosphorous and potassium fertility regimes affect tomato transplant growth. *Hort Sci.*, **26**(2): 141-142.
- Midan, A.A., N. M. Malash and M. M. El-ssayed. 1998. Intensification and nitrogen fertilization in relation to tomato yield. *Ann. Agric. Sci. Fac. Agric. Ainshams Univ., Cairo, Egypt*. **30**(2): 1413-1431.

- Mohanty, B. and K. and Hossain, M. M. 2001. Performance of hybrid tomato cv. Vaishali to nitrogen, potash and micronutrient application. *Advances in Plant Sciences*. **14**(1):185-190.
- Mukhopadhyay, D., Eunus, M., and Haque, M. M. 1986. Response of major crop to balanced fertilizer application. D.A.E and FAO, Publ. Field Document. 5, 1p.
- Nanadal. J. K., Ramesh, V., Pandey, U. C. and Vasist, R. 1998. Effect of phosphorus and potassium on growth yield and quality of tomato. *J. Potassium Res.*, **14**(1-4): 44-49.
- Nassar, H. H. 1988. Effect of planting pattern, plant population and nitrogen level on yield and quality of tomato. *Acta Hort.*, **190**: 435-442.
- Olsen, S. R., Cole, C. V., Watanabe, F. S. and Dean, L. A. 1965. Estimation of available phosphorous in soil by extraction with sodium bicarbonate. U. S Dept. Agric. Cire. p. 929.
- Page, A. L., Miller, R. H. and Keency, D. R. 1982. Methods of Soil Analysis. Part II (2nd ed.). Amer. Soc. Agron. Inc. Madison, Winsconsin, USA.
- Pandey, R.P., P.N. Solanki, R.K. Saraf and M.S. Parihar. 1996. Effect of nitrogen and phosphorus on growth and yield of tomato (*Lycopersicon esculentum* Mill.) varieties. *Punjab Veg. Grow*, **32**: 1-5.
- Pansare, P.D., B.B. Desai and U.D. Chavan. 1994. Effects of different nitrogen, phosphorus and potassium ratios on yield and quality of tomato. *J. Maharashtra Agric. Univ., India.*, **19**(3): 462-463.
- Patil, A. A. and Bojoppa, K. M. 1984. Effects of cultivars and graded levels of nitrogen and phosphorous on certain quality attributes of tomato (*Lycopersicon esculentum* Mill.). II sugars, dry matter content and juice percentage. *Mysore J. Agril. Sci.*, **18**(4): 292-295.
- Piper, C.S. 1942. Soil and plant analysis. Adelaide Univ. press. Australia.
- Rashid, M. A. 1993. Sabji Biggan. Bangla Accademi, Dhaka, Bangladesh. pp. 234-269.

- Razia, S. and M.S. Islam. 1980. The effect of phosphatic fertilizer on the yield of tomato. *Bangladesh Hort.*, **8**(2): 29-31.
- Rhoads, F. M., Gardner, C. S., Mbuya, O. S., Queeley, G. L. and Edwards, H. M. 1999. Tomato fertilization, ground cover, and soil nitrate nitrogen movement. *Proceedings Florida State Horticultural Society*. **2000**: (1-2). 315-319.
- Sainju, U. M., Singh, B. P. and Whitehead, W. F. 2000. Cover crops and nitrogen fertilization effects on soil carbon and nitrogen and tomato yield. *Canadian J. Soil Sci.*, **80**(3): 523-532.
- Sainju, U. M., Syed, R. and Singh B. P. 2001. Evaluating hairy vetch residue as nitrogen fertilizer for tomato in soilless medium. *Hort. Sci.*, **36**(1): 90-93.
- Salunkhe, D. K., Desai, B. B. and N. R. Bhat. 1987. Vegetable and flower seed production. 1st Edn. Agricola Publishing Academy, New Delhi, India. pp. 118-119.
- Sandoval, V. M., Guertal, E. A. and Wood, C. W. 2001. Greenhouse tomato response to low ammonium-nitrogen concentrations and duration of ammonium nitrogen supply. *J. Plant Nutri.*, **24**: 11.
- Scholberg, J., McNeal, B L., Boote, K. J., Jones, J. W., Locascio, S. J. and Olson, S. M. 2000. Nitrogen stress effects on growth and nitrogen accumulation by field-grown tomato. *Agronomy J.*, **92**(1): 159-167.
- Seno, S., Nakagawa, J., Zanin, A. C. W. and Mischon, M. M. 1987. Effects of phosphorous and potassium levels on fruits characteristics and quality of tomato seeds. *Hort. Brasileira*, **5**(2): 25-28.
- Sharma, C. B. and Mann, H. S. 1971. Relative response of phosphatic fertilizers at varying levels of nitrogen and phosphorous in tomato. *Indian J. Hort.*, **28**(1): 46-54.
- Sharma, K. C., Singh, A. K. and Sharma. S. K. 1999. Studies on nitrogen and phosphorus requirements of tomato hybrids. *Annals of Agricultural Research*. **20**(4): 399-402.


- Silva, F. C., Miranda, J. R. P., and Alvarenga, M. A. R. 2001. Yield and nutrient concentration of tomato plants pruned and grown under high planting density in relation to phosphorus, gypsum and nitrogen sources. *Horticultura Brasileira*. **19**(1):64-69.
- Singh, A. K. Singh, P. K. and Gaur, G. S. 2000. Determination of nitrogen doses and its method of application for growth and yield of tomato (*Lycopersicon esculentum* Mill.) var. Pusa hybrid-2. *Haryana J. Hort. Sci.*, **29**(3-4), 263-264.
- Singh, K. and Kumer, S. 1969. Effect of nitrogen and phosphorous fertilization on the growth and yield of onion (*Allium cepa* L). *J. Res. Ludhiana.*, **6**: 764-768.
- Singh, T. R., Singh, S., Singh, S. K., Singh, M. P. and Srivastava, B. K. 2004. Effect of integrated nutrient management on crop nutrient uptake and yield under okra-pea-tomato cropping system in a Mollisol. *Indian J. Hort.*, **61**(4): 312-314.
- Song, K., Jiao, X., Li, L., Yan, J., Song, K. M., Jiao, X. Z., Li, L. and Yan, J. Q. 1998. Effects of phosphate starvation on growth state and phosphorus uptake of tomato seedlings. *Acta Botanica Yunnanica*. **20**(3): 343-350.
- Soares, K., T. Kimoto, F.A.D. Conceicao, La De Lima, C. Kurozawa, J. Nakagawa, and A. E. Boaretto. 1985. Mineral fertilization of tomato (cv. Petomech). II. Effect of nitrogen fertilization on a red yellow latosol sandy phase. Faculdade de Ciencias Agronomicas, UNESP, 18600 Botucatu, SP, Brazil. *Cientifica*. **13**(1/2) :51-57.
- Taya, J. S., Malik, Y. S. Pandita, M. L. and Khurana, S. C. 1994. Fertilizer management in potato based cropping system II: effect of residual and applied fertility on first succeeding crops of tomato and okra. *J. Indian Potato Association.*, **21**(3/4): 189-194.
- Topeuoglu, B. and S. R Yalcin. 1994. Effect of lime and phosphorus on the formation of oxalic acid and on the yield in tomato. Ankara Universities Ziraat Fakultesi Yilligi, **44**(1/2): 171-181.



- Trpevski, V. Demirovska, V. Stojcevska, L. and Oggenovski, V. 1992. The effect of different rates of nitrogen on the quality and yield of some tomato cultivars. *Savremena Poljoprivreda*. **40**: (1-2): 100- 105.
- UNDP. 1988. Land Resource Appraisal of Bangladesh for Agricultural Development Report 2: Agro-ecological Regions of Bangladesh. FAO, Rome, Italy, pp. 577.
- Valenzuela, J.L., A. Sanchez, and L. Romero. 1987. Ascending critical nitrogen supply by means of nitrate reductase activity in tomato and cucumber plants of Plant nutrition. **10**(9/16):1733-1741.
- Wateren, D.R. and Coltman, R.R. 1988. Phosphorus concentration and application interval influence growth and mycorrhizal infection of tomato and onion transplants. *J. American Soc. Hort. Sci.*, **113**(5): 704-708.
- Winsor, G.W. and M.I.E. Long. 1962. The effect of nitrogen, phosphorus, potassium, magnesium and lime in factorial combination on ripening disorders of glasshouse tomatoes. *J. Hort. Sci.*, **42**:391-402.
- Xin, X. Y., Hui, L. J., and Lili, H. 1997. The effect of N, P, K mixed application on yields and quality of tomato in Solar green house. *China Veg.*, **4**:10-13.

APPENDICES

Appendix I: Map showing the experimental site under the study



Appendix II: Monthly record of air temperature, relative humidity and sunshine of the experimental site during the period from November 2008 to February 2009

Year	Month	Air Temperature (°C)		Relative Humidity (%)	Rainfall (mm)	Sunshine (hr)
		Maximum	Minimum			
2008	October	30.4	15.6	81	0	7.3
	November	32.3	16.3	79	0	7.9
	December	29.0	13.0	72	1	3.9
2009	January	28.1	11.1	55	1	5.7
	February	33.9	12.2	67	45	8.7
	March	34.6	16.5	65	88	7.3

Source: Bangladesh Meteorological Department (Climate & water division)
Agargoan, Dhaka- 1207

Appendix III: Analysis of variance of the data on plant height at 30, 60 DAT, at harvest, number of leaves per plant and 60 DAT and number of branches per plant at harvesting period as influenced by nitrogen and phosphorous

Source of variation	Degrees of freedom	Mean square				
		Plant height at			No. of leaves per plant at	No. of branches per plant at
		30 DAT	60 DAT	Harvest	60 DAT	Harvest
Replication	2	0.250	0.813	1.188	2.313	0.015
Nitrogen(A)	3	40.438*	130.779**	196.545*	319.071**	5.676**
Phosphorous (B)	3	40.107*	78.760**	171.048*	173.876**	9.174**
Interaction (A×B)	9	1.290*	1.573*	0.647*	8.359*	0.151*
Error	30	3.850	4.479	1.388	6.179	0.226

** 1% level of significance

* 5% level of significance

NS: Non Significance

Appendix IV: Analysis of variance of the data on number of cluster / plant, number of flowers/ cluster, number of flowers/ plant, number of fruits/ cluster, number of fruits/ plant influenced by nitrogen and phosphorous

Source of variation	Degrees of freedom	Mean square				
		Number of cluster / plant	Number of flowers/ cluster	Number of flowers/ plant	Number of fruits/ cluster	Number of fruits/ plant
Replication	2	0.010	0.174	0.348	0.021	7.563
Nitrogen (A)	3	14.279**	0.466**	489.450**	0.668**	351.094**
Phosphorous (B)	3	12.892**	0.734**	505.157**	0.159 ^{NS}	274.154**
Interaction (A×B)	9	0.185*	0.021*	8.442*	0.018*	3.966*
Error	30	0.142	0.038	5.704	0.061	4.096

** 1% level of significance

* 5% level of significance

NS: Non Significance

Appendix V: Analysis of variance of the data on fruit size, weight of individual fruit (gm), fruit weight/plant (kg) and fruit and fruit yield (ton/ha) as influenced by nitrogen and phosphorous.

Source of variation	Degrees of freedom	Mean square				
		Fruit size		Weight of individual fruit (gm)	Fruit weight /plant (kg)	Fruit yield (ton/ha)
		Length (cm)	Breadth (cm)			
Replication	2	0.006	0.010	4.938	0.011	1.563
Nitrogen (A)	3	1.361**	0.270**	819.407**	2.703**	3515.749**
Phosphorous (B)	3	0.781**	0.139**	262.031**	1.701**	2208.871**
Interaction (A×B)	9	0.043*	0.006*	14.812*	0.051**	65.709**
Error	30	0.019	0.011	6.471	0.005	5.296

** 1% level of significance

* 5% level of significance

NS: Non Significance



Appendix VI: Analysis of variance of the data on pH of post harvest soil, nitrogen (%), phosphorous (ppm) of post harvest soil as influenced by nitrogen and phosphorous.

Source of variation	Degrees of freedom	Mean square		
		pH	Nitrogen (%)	Phosphorous (P) in ppm
Replication	2	0.010	0.000	5.290
Nitrogen (A)	3	0.072 ^{NS}	0.001*	218.220**
Phosphorous (B)	3	0.092 ^{NS}	0.000*	907.005**
Interaction (A×B)	9	0.097 ^{NS}	0.000 ^{NS}	11.762*
Error	30	0.038	0.000	4.117

** 1% level of significance * 5% level of significance NS: Non Significance

Appendix VII: Analysis of variance of the data on nitrogen (%) in shoot, phosphorous (P) in ppm in shoot, N % in fruit, P % in fruit as influenced by nitrogen and phosphorous

Source of variation	Degrees of freedom	Mean square			
		Nitrogen (%) in straw	Phosphorous (P) in ppm in straw	N in fruit (%)	P in fruit (%)
Replication	2	0.001	3.755	0.109	0.000
Nitrogen (A)	3	2.175**	1346.497**	1.664**	0.005**
Phosphorous (B)	3	0.337**	1761.885**	0.299**	0.010**
Interaction (A×B)	9	0.026*	79.166**	0.013*	0.000*
Error	30	0.008	5.455	0.025	0.000

** 1% level of significance * 5% level of significance NS: Non Significance

Shere-Bangla Agricultural University
Library

Accession No. 38728
Sign: _____ Date: 24.2.15

স্বাক্ষরিত ও সত্যায়িত
সংগ্রহ নং 55(06) সাই.সি.
তারিখ: 9/08/11